Mathematical Institute
Slovak Academy of Sciences, Bratislava

Preprint 14/1996

Anatolij Dvurečenskij
Karl Svozil

Product of partition logics, orthoalgebras and automata

June 12, 1996

PREPRINT SERIES

Product of Partition Logics, Orthoalgebras and Automata

Anatolij DVUREČENSKIJ[1], Karl SVOZIL[2]

[1] Mathematical Institute, Slovak Academy of Sciences,
Štefánikova 49, SK-814 73 Bratislava, Slovakia
e-mail: dvurecen@mau.savba.sk
[2]Institute for Theoretical Physics, Technical University of Vienna
Wiedner Hauptstr. 8–10/136, A-1040 Vienna, Austria
e-mail: svozil@tph.tuwien.ac.at

Abstract

We present an attempt how to define a coupled system consisting of two partitions logics and we introduce a product of partition logics. These partition logics have a close connection with Moore and Mealy type automata. We show how the coupled system of two automata is connected with the product of partition logics, and some illustrative examples are presented.

1 Introduction

In the axiomatic approach to quantum mechanics, the event structure of a physical system is identified with a quantum logic [3] or an orthoalgebra [21, 9] while in the case of a classical mechanics with a Boolean algebra [20]. Assume that we have two independent physical systems with event structures P and Q, and we wish to regard them as a coupled system. The event structure L of this coupled system is usually called a tensor product of P and Q and we write $L = P \otimes Q$.

Tensor products in various approaches have been studied in [21, 1, 17, 7, 11, 15, 16, 10, 26, 27]. A tensor product of orthoalgebras has been investigated by Foulis and Bennett in [8] via a universal mapping property, and a tensor product of an orthoalgebra and a Boolean algebra is given in [10].

A little more general structure than orthoalgebras are quasi orthoalgebras for which a binary operation $\oplus$ is not, in general, associative. A very important example of quasi orthoalgebras, partition logics, have an intimate connection [6] with the investigation of the empirical propositional structure of Moore and Mealy type automata.

The aim of the present paper is to introduce a weaker form of a tensor product of partition logics, a product of partition logics, and to show its connection with automata. We show that this structure exists.

2 Orthoalgebras

The notion of orthoalgebras (or quasi orthoalgebras) goes back to axiomatic models of quantum mechanics introduced by Foulis and Randall [11, 21] as special algebraic structures describing propositional logics.

[1]The paper has been partially supported by the grant G 229/94 SAV, Bratislava, Slovakia, and by the Mitteln zur Förderung der Auslandsbeziehungen an der Technischen Universität Wien.

A *quasi orthoalgebra* is a set L endowed with two special elements $0, 1 \in L$ $(0 \neq 1)$ and equipped with a partially defined binary operation $\oplus$ satisfying the following conditions for all $a, b \in L$:

(oai) if $a \oplus b$ is defined, then $b \oplus a$ is defined and $a \oplus b = b \oplus a$ (commutativity law);
(oaii) $a \oplus 0$ is defined for any $a \in L$ and $a \oplus 0 = a$;
(oaiii) for any $a \in L$, there is an element $a' \in L$ such that $a \oplus a'$ is defined and $a \oplus a' = 1$ (orthocomplementation law);
(oaiv) if $a \oplus (a' \oplus b)$ is defined, then $b = 0$;
(oav) if $a \oplus (a \oplus b)$ is defined, then $a = 0$;
(oavi) if $a \oplus b$ is defined, then $a \oplus (a \oplus b)'$ is defined and $b' = a \oplus (a \oplus b)'$.

The following facts are true:

Proposition 2.1 *Let L be a quasi orthoalgebra, $a, b \in L$. Then*

(a) $0' = 1$, $1' = 0$;
(b) $(a')' = a$;
(c) *if* $a \oplus b = a \oplus c$, *then* $b = c$;
(d) *if* $a \oplus b = 1$, *then* $b = a'$.

The unique element a' is called *orthocomplement* of $a \in L$, and the unary operation $' : L \to L$ defined by $a \mapsto a'$, $a \in L$, is said to be an *orthocomplementation*. We shall say that two elements $a, b \in L$ (i) are *orthogonal*, and write $a \perp b$, iff $a \oplus b$ is defined in L (it is clear that $a \perp b$ iff $b \perp a$), and (ii) $a \leq b$ iff there is an element $c \in L$ with $a \oplus c = b$.

It is easily to shown that the relation $\leq$ is reflexive and antisymmetric, but needs not to be transitive. An associative quasi orthoalgebra, i.e., a quasi orthoalgebra, for which the associative law

(oavii) if $a \oplus b$, $(a \oplus b) \oplus c$ are defined in L, so are $b \oplus c$ and $a \oplus (b \oplus c)$, and $(a \oplus b) \oplus c = a \oplus (b \oplus c)$

holds is said to be an *orthoalgebra* (OA in abbreviation). If in L there are the elements $(a \oplus b) \oplus c$, and $a \oplus b(\oplus c)$ and if they coincide, we denote it as $a \oplus b \oplus c$. In any orthoalgebra, $\leq$ is transitive. On other hand it is possible to give an example of a quasi orthoalgebra with transitive $\leq$ which does not correspond to any orthoalgebra. If the elements $a \oplus (b \oplus c)$ and $(a \oplus b) \oplus c$ exist in a quasi orthoalgebra and if they coincide, we denote them as $a \oplus b \oplus c$.

Due to Golfin [12], an orthoalgebra is a set L with two special elements $0, 1 \in L$ $(0 \neq 1)$ and endowed with a partial binary operation $\oplus$ satisfying (oai), (oaii), (oaiii), (oavii), and (oav*) if $a \oplus a$ is defined, then $a = 0$.

The original idea of the partial binary operation $\oplus$ goes back to Boole's pioneering paper [2], where he wrote $a + b$ as the logical disjunction of events a and b when the logical conjuction $ab = 0$, so that, for mutually excluding events a and b, $a + b$ is defined. This is all that is needed for probability theory: if $ab = 0$, then $P(a + b) = P(a) + P(b)$. To avoid confusion, we write $a \oplus b$ for $a + b$ when $ab = 0$.

Note that one can rewrite axioms for a Boolean algebra in terms of Boole's ideas of $a + b$. For more details, see Foulis and Bennett [8].

In addition, let L be an orthomodular poset (OMP for abbreviation) (or an orthomodular lattice, OML in short), i.e., a poset L with the least and last elements 0 and 1 and a unary operation $^{\perp} : L \to L$, called an *orthocomplementation*, such that, for all $a, b \in L$,

(i) $(a^{\perp})^{\perp} = a$;

(ii) if $a \leq b$, then $b^{\perp} \leq a^{\perp}$;
(iii) $a \vee a^{\perp} = 1$;
(iv) if $a \leq b^{\perp}$ (and we write $a \perp b$), then $a \vee b \in L$;
(v) if $a \leq b$, then $b = a \vee (a \vee b^{\perp})^{\perp}$.

(For OML, L has to be additionally a lattice). Then L can be organized into an OA if the binary operation $\oplus$ is defined via $a \oplus b$ exists in L iff $a \leq b^{\perp}$ and $a \oplus b := a \vee b$. The unary operation $' : L \to L$ is defined via $a' := a^{\perp}$, $a \in L$.

We recall that if L is an OA and $a, b \in L$ are mutually orthogonal, then $a, b \leq a \oplus b$, and $a \oplus b$ is the minimal upper bound for a and b (i.e., $a, b \leq a \oplus b$, and if there is $c \in L$ with $a, b \leq c \leq a \oplus b$, then $c = a \oplus b$), but this does not mean that $a \vee b$ exists in L, so that L cannot be necessarily an OMP.

A subset A of a quasi OA (OA) L is a *quasi suborthoalgebra* (*suborthoalgebra*) of L is (i) $0, 1 \in A$; (ii) if $a \in A$, then $a' \in A$; (iii) $a, b \in A$ with $a \perp b$ implies $a \oplus b \in L$.

If a (quasi) suborthoalgebra A of L is, in addition, a Boolean algebra with respect to $\leq$, A is called a *Boolean suborthoalgebra* of L. Denote by $\vee_A$ and $\wedge_A$ the join and the meet taken only in A, respectively. Then, $a \oplus b = a \vee_A b$ whenever $a, b \in A$ and L is an OA. A maximal Boolean suborthoalgebra of L is called a *block*.

3 Partition Logics

In this section, we present a notion of partition logics which will have an intimate connection with special types of automata, and which will generalize the results of Svozil [25] and Schaller and Svozil [22, 23, 24].

Let L be a quasi orthoalgebra with $\leq$. A non-void subset I of L is said to be an *ideal* of L if

(i) if $a \in I, b \in L, b \leq a$, then $b \in I$;
(ii) $a, b \in I$ with $a \perp b$ imply $a \oplus b \in I$.

It is clear that $0 \in I$. An ideal I of L is said to be (i) *proper* if $I \neq L$ or, equivalently, $1 \notin I$; (ii) *prime* if, for any $a \in L$, either $a \in I$ or $a' \in I$. We denote by $P(L)$ the set of all prime ideals in L.

A *probability measure* (or also a *state*) on L is a mapping $s : L \to [0, 1]$ such that (i) $s(1) = 1$, and (ii) $s(a \oplus b) = s(a) + s(b)$ whenever $a \perp b$. A probability measure s is *two-valued* if $s(a) \in \{0, 1\}$ for any $a \in L$.

We recall that there is a one-to-one correspondence between two-valued probability measures and prime ideals: If s is a two-valued probability measure, then $I_s = \{a \in L : s(a) = 0\}$ is a prime ideal; and if I is a prime ideal, then $s_I : L \to [0, 1]$ defined via $s_I(a) = 0$ iff $a \in I$, otherwise $s_I(a) = 1$, is a two-valued probability measure on L.

A set $\mathcal{S}$ of probability measures on L is called *separating* if for all $a, b \in L$, $a \neq b$, there is a probability measure $s \in \mathcal{S}$ such that $s(a) \neq s(b)$. L is called *prime* iff it has a separating set of two-valued probability measure or, equivalently, for any different elements $a, b \in L$ there is a prime ideal I of L such that $a \in I$ and $b \notin I$.

Let $\mathcal{L}$ be a family of of quasi orthoalgebras (or OAs, OMP, Boolean algebras, etc.) satisfying the following conditions: For all $P, Q \in \mathcal{L}$, $P \cap Q$ is a quasi suborthoalgebra (subOA, sub OMP, Boolean subalgebra, etc.) of both P and Q, and the partial orderings and orthocomplementations coincide on $P \cap Q$. Define the set $L = \bigcup := \bigcup\{P : P \in \mathcal{L}\}$, a relation $\oplus$ and the unary operation $'$ as follows:

(i) $a \oplus b$ iff there is a $P \in \mathcal{L}$ such that $a, b \in P$ and $a \perp_P b$, then $a \oplus b = a \oplus_P b$;
(ii) $a' = b$ iff there is a $P \in \mathcal{P}$ such that $a, b \in P$ and $a^{'_P} = b$.



Figure

Figure 1: The Fano plane.

The set L with the above defined $\oplus$ is called the *pasting* of the family $\mathcal{L}$.

Let $\mathcal{R}$ be a family of finite partitions of a fixed set X. The pasting of the family of Boolean algebras $\{B_R : R \in \mathcal{R}\}$ is called *partition logic*, and we denote it as a couple $(X, \mathcal{R})$.

We notify that, for $a, b \in L = (X, \mathcal{R})$, $a \oplus b$ is defined on L iff there exists a decomposition $R \in \mathcal{R}$ such that $a, b \in B(R)$ and $a \cap b = \emptyset$, where $B(R)$ is a Boolean algebra generated by R; then $a \oplus b := a \cup b$.

We recall that two quasi orthoalgebras L_1 and L_2 are *isomorphic* iff there is a one-to-one mapping $\phi : L_1 \to L_2$ such that $a \oplus b$ is defined in L_1 iff $\phi(a) \oplus \phi(b)$ is defined in L_2 and $\phi(a \oplus b) = \phi(a) \oplus \phi(b)$.

The following result [6] describes quasi orthoalgebras isomorphic to partition logics.

Theorem 3.1 *A quasi orthoalgebra L is isomorphic to a partition logic if and only if L is prime.*

It is worth no notify that if in a prime quasi orthoalgebra L there exist elements $x := (a \oplus b) \oplus c$ and $y := a \oplus (b \oplus c)$, for $a, b, c \in L$, then $x = y$. Indeed, by Theorem 3.1, there is a separating system of two-valued probability measures on L, $\mathcal{S}$, and a probability measure $s \in \mathcal{S}$ such that $s(x) \neq s(y)$. Then $s(a \oplus b) + s(c) \neq s(a) + s(b \oplus c)$ which gives the contradiction $s(a) + s(b) + s(c) \neq s(a) + s(b) + s(c)$.

In any rate, the existence of one of x or y in L does not imply the existence of the second one in L, see Example 4.1 below.

4 Coupled Systems of Partition Logics

The tensor product of orthoalgebras in the category of orthoalgebras has been studied by Foulis and Bennett [8]. They showed that if both orthoalgebras P and Q have "enough" probability measures, then the tensor product of P and Q exists. However, they have found an example of orthoalgebra, the Fano plane drawn in Fig . 1, for which the tensor product $P \otimes P$ fails in the category of orthoalgebras. Dvurečenskij [4] showed that if we use a more general structure, effect algebras, then the tensor product of the Fano plane with itself exists in the category of effect algebras. In general, if both effect algebras P and Q have a non-empty system of probability measures, then the tensor product $P \otimes Q$ exists in the category of effect algebras.

Partition logics have a separating system of two-valued probability measures, unfortunately, they are not, in general, effect algebras, because they are not even (associative) orthoalgebras. Therefore, for the tensor product of partition logics we have to look for another way how to introduce it.

4.1 Product of Partition Logics

Assume that $(X, \mathcal{R})$ and $(Y, \mathcal{S})$ be two partition logics, where $\mathcal{R}$ and $\mathcal{S}$ are two systems of finite decompositions of X and Y, respectively. Choose $R \in \mathcal{R}$ and $S \in \mathcal{S}$. Then $R \times S := \{A \times B : A \in R, \ B \in S\}$ is a decomposition of $X \times Y$. Define

$$\mathcal{R} \times \mathcal{S} := \{R \times S : R \in \mathcal{R}, S \in \mathcal{S}\}.$$

Then the partition logic $\{X \times Y, \mathcal{R} \times \mathcal{S}\}$ is called the *product partition logic* of $(X, \mathcal{R})$ and $(Y, \mathcal{S})$.

Example 4.1 *Let $\Omega = \{1, 2, 3, 4, 5, 6\}$ and let B_1 and B_2 be the Boolean algebras generated by $R_1 := \{\{1\}, \{2\}, \{3\}, \{4\}, \{5, 6\}\}$ and $R_2 : \{\{1\}, \{2\}, \{3, 4\}, \{5\}, \{6\}\}$, respectively. The system $(\Omega, \{R_1, R_2\})$ is a partition logic which is not a Boolean algebra, it is only a quasi orthoalgebra. Then $(\Omega \times \Omega, \{R_1 \times R_1, R_1 \times R_2, R_2 \times R_1, R_2 \times R_2\})$ is a product partition logic of $(\Omega, \{R_1, R_2\}\}$ with itself.*

Proposition 4.2 *Let μ and ν be probability measures on the partition logics $(X, \mathcal{R})$ and $(Y, \mathcal{S})$, respectively, where card $(Y) = n$ and $\mathcal{S} = \{\{y_1\}, \ldots, \{y_n\}\}$, $y_i \in Y$ for $i = 1, \ldots, n$, $y_i \neq y_j$ for $i \neq j$, and $n \geq 1$. Then there is a unique probability measure $\mu \times \nu$ on $(X \times Y, \mathcal{R} \times \mathcal{S})$ such that*

$$(\mu \times \nu)(A \times B) = \mu(A)\, \nu(B), \ \ A \in (X, \mathcal{R}), \ \ B \in (Y, \mathcal{S}). \tag{1}$$

If, in addition, μ and ν are two-valued measures, so is $\mu \times \nu$.

Proof. Let U be any element of the product partition logic $L = (X \times Y, \mathcal{R} \times \mathcal{S})$. Without loss of generality we can suppose that $Y = \{1, \ldots, n\}$. It is easy to show that U can be represented uniquely in the form $U = \bigcup_{i=1}^n A_i \times \{i\}$, where all A_i's belong to the same Boolean algebra B_R generated by a decomposition $R \in \mathcal{R}$; the case $A_i = \emptyset$ is not excluded.

In addition, $U := \bigcup_{i=1}^n A_i \times \{i\} \perp \bigcup_{i=1}^n B_i \times \{i\} =: V$ iff $\{A_i\}_{i=1}^n$ and $\{B_i\}_{i=1}^n$ belong to the same Boolean algebra B_R and $A_i \cap B_i = \emptyset$ for $i = 1, \ldots, n$, and in this case $\bigcup_{i=1}^n A_i \times \{i\} \oplus \bigcup_{i=1}^n B_i \times \{i\} = \bigcup_{i=1}^n (A_i \cup B_i) \times \{i\} = \bigcup_{i=1}^n (A_i \oplus B_i) \times \{i\}$.

If now we define $(\mu \times \nu)(U) := \sum_{i=1}^n \mu(A_i)\nu(\{i\})$, then $\mu \times \nu$ is a well-defined mapping on L. It is simple to verify that $X \times Y = \bigcup_{i=1}^n X \times \{i\}$. Hence $(\mu \times \nu)(1) = (\mu \times \nu)(X \times Y) = \sum_{i=1}^n \mu(X)\nu(\{i\}) = \mu(X)\nu(Y) = 1$.

Similarly, $(\mu \times \nu)(U \oplus V) = (\mu \times \nu)(\bigcup_{i=1}^n (A_i \oplus B_i) \times \{i\}) = \sum_{i=1}^n \mu(A_i \oplus B_i)\nu(\{i\}) = \sum_{i=1}^n (\mu(A_i) + \mu(B_i))\nu(\{i\}) = (\mu \times \nu)(U) + (\mu \times \nu)(V)$, so that $\mu \times \nu$ is a probability measure on L satisfying (1). The uniqueness of $\mu \times \nu$ is now evident. $\square$

The probability measure $\mu \times \nu$, if it exists, is said to be a *product probability measure* of μ and ν. We note that the extension of $\mu \times \nu$ from (1) to a probability measure on the product of general partition logics seems to be open.

4.2 Tensor Product of Prime Quasi Orthoalgebras

According to Theorem 3.1, any prime quasi orthoalgebra is isomorphic to a partition logic. Therefore, the coupled system consisting of two prime orthoalgebras P and Q being again a prime quasi orthoalgebra can be called a tensor product of P and Q and it is denoted as $P \otimes Q$. In addition, they have an intimate connection with the product of isomorphic partition logics. For a rigorous introduction of the tensor product we need the following definitions.

Let P and L be two quasi orthoalgebras. A mapping $\phi : P \to L$ is said to be

(i) a *morphism* iff $\phi(1) = 1$, and $p \perp q$, $p, q \in P$, implies $\phi(p) \perp \phi(q)$ and $\phi(p \oplus q) = \phi(p) \oplus \phi(q)$;

(ii) a *monomorphism* iff ϕ is a morphism and $\phi(p) \perp \phi(q)$ iff $p \perp q$;

(iii) an *isomorphism* iff ϕ is a surjective monomorphism.

Let P, Q, L be quasi orthoalgebras. A mapping $\beta : P \times Q \to L$ is called a *bimorphism* iff

(i) $a, b \in P$ with $a \perp b$, $q \in Q$ imply $\beta(a, q) \perp \beta(b, q)$ and $\beta(a \oplus b, q) = \beta(a, q) \oplus \beta(b, q)$;

(ii) $c, d \in Q$ with $c \perp d$, $p \in P$ imply $\beta(p, c) \perp \beta(p, d)$ and $\beta(p, c \oplus d) = \beta(p, c) \oplus \beta(p, d)$;

(iii) $\beta(1, 1) = 1$.

If $\beta : P \times Q \to L$ is a bimorphism, then $\beta(\cdot, 1) : P \to L$ and $\beta(1, \cdot) : Q \to L$ are morphisms. Therefore, for $p \in P$ and $q \in Q$, we have $\beta(p, 1)^\perp = \beta(p^\perp, 1)$, $\beta(1, q)^\perp = \beta(1, q^\perp)$ and $\beta(p, 0) = \beta(0, q) = 0$.

Also, if $a, b, p \in P$ and $c, d, q \in Q$, we have $a \leq b \Rightarrow \beta(a, q) \leq \beta(b, q)$ and $c \leq d \Rightarrow \beta(p, c) \leq \beta(p, d)$.

Definition 4.3 *Let P and Q be prime quasi orthoalgebras. We say that a pair (T, τ) consisting of a prime quasi orthoalgebra T and a bimorphism $\tau : P \times Q \to T$ is a tensor product of P and Q iff the following conditions are satisfied:*

(i) *If L is a quasi orthoalgebra and $\beta : P \times Q \to L$ is a bimorphism, there exists a morphism $\phi : T \to L$ such that $\beta = \phi \circ \tau$.*

(ii) *For every element of $t \in T$, there is a block B of T such that $t = \bigoplus_{i=1}^n \tau(p_i, q_i)$, where $\tau(p_i, q_i)$ belongs to the block B for $i = 1, \dots, n$.*

It is not hard to show that if a tensor product (T, τ) of P and Q exists, it is unique up to an isomorphism, i.e., if (T, τ) and (T^*, τ^*) are tensor products of D-posets P and Q, then there is a unique isomorphism $\phi : T \to T^*$ such that $\phi(\tau(p, q)) = \tau^*(p, q)$ for all $p \in P$, $q \in Q$. Unless confusion threatens, we usually refer to $P \otimes Q$ rather than to $(P \otimes Q, \otimes)$, as being a tensor product.

Unfortunately, the authors do not know the conditions under those the tensor product of prime quasi orthoalgebras exists in the category of prime quasi orthoalgebras.

If both components P and Q are prime orthoalgebras, then the tensor product of P and Q exists in the category of orthoalgebras. This follows from Theorem 6.1 [8]. Indeed, if $p \in P$ and $q \in Q$ are two non-zero elements of the prime orthoalgebras P and Q, then there exist two probability measures μ and ν on P and Q such that $\mu(p) = 1 = \nu(q)$ (if $p = 1$ then $\mu(1) = 1$ for any μ, if $p \neq 1$ then $0 \neq p^\perp \neq 1$ and the primeness and the separateness of P entail the existence of a two-valued probability measure μ with $0 = \mu(p^\perp) \neq \mu(1) = 1$). By Theorem 6.1 of Foulis and Bennett, this is a sufficient condition for the existence of the tensor product of P and Q in the category of orthoalgebras. In addition, for all probability measures μ and ν on P and Q, respectively, the product measure $\mu \times \nu$, defined by $\mu \times \nu(p \otimes q) = \mu(p)\nu(q)$, $p \in P$, $q \in Q$, exists. However, we do not know whether this tensor product exists also in the category of prime quasi orthoalgebras.

More hopeful seems to be the coupled system consisting of a prime quasi orthoalgebra and a Boolean algebra, called a bounded Boolean power of a prime quasi orthoalgebra, see also similar problems in [4] for D-posets and [10] for orthoalgebras. The authors hope to present some results for these coupled systems in the future.

5 Realization by Automata

The product of finite automata logics has an intuitive and rather simple realization: If the sets of states of two automata $M_1 = (S_1, I_1, O_1, \delta_1, \lambda_1)$ and $M_2 = (S_2, I_2, O_2, \delta_2, \lambda_2)$ are mutually disjunct, then the automaton partition logic of the *parallel decomposition* $M_1 \| M_2$ of the two automata is the product $\{S_1 \times S_2, \mathcal{R}_1 \times \mathcal{R}_2\}$ of the automaton partition logics $\{S_1, \mathcal{R}_1\}$ and $\{S_2, \mathcal{R}_2\}$ associated with M_1 and M_2, respectively. For a definition of the notation, see below.

5.1 Moore and Mealy Automata

A *finite sequential machine* or *automaton* is a device with the following properties [18, 14, 13]: (i) A finite set of inputs which can be applied in a sequential order; (ii) a finite set of internal configurations or states; (iii) a finite set of outputs; (iv) the present internal configuration and input uniquely determine the next internal configuration and the output.

A *Moore (Mealy) automaton* is a quintuple $M = (S, I, O, \delta, \lambda)$, where

(i) S is a finite (nonempty) set of states;

(ii) I is a finite (nonempty) set of inputs;

(iii) O is a finite (nonempty) set of outputs;

(iv) $\delta : S \times I \longrightarrow S$ is a computable transition function;

(v) $\lambda : S \longrightarrow O$ is a computable output function (Moore automaton);

(v') $\lambda : S \times O \longrightarrow O$ is a computable output function (Mealy automaton).

In what follows and if not mentioned otherwise, s, i, o stand for a particular internal state, input and output, respectively. Moore (Mealy) machines are represented by flow tables and state graphs.

Example 5.1 *To illustrate this, assume a Mealy machine $M_s = (S, I, O, \delta, \lambda)$ which has n states, n inputs and 2 outputs. That is*

$$\begin{aligned} S &= \{1, 2, ..., n\}, \\ I &= \{1, 2, ..., n\}, \\ O &= \{0, 1\}. \end{aligned}$$

Its transition and output functions are ($\delta_{s,x}$ stands for the Kronecker delta function)

$$\begin{aligned} \delta(s, i) &= i, \\ \lambda(s, i) &= \delta_{s,i} = \begin{cases} 1 & \text{if } s = i \\ 0 & \text{if } s \neq i \end{cases} \end{aligned}$$

The flow table and state graph (for n=3) of this Mealy automaton is given in Fig. 2.

5.2 Machine Isomorphism, Serial and Parallel Decompositions

Two automata $M_1 = (S_1, I_1, O_1, \delta_1, \lambda_1)$ and $M_2 = (S_2, I_2, O_2, \delta_2, \lambda_2)$ of the same type are *isomorphic* if and only if there exist three one-to-one mappings $f : S_1 \longrightarrow S_2$, $g : I_1 \longrightarrow I_2$, $h : O_1 \longrightarrow O_2$ such that $f[\delta_1(s_1, i_1)] = \delta_2[f(s_1), g(i_1)]$ and

$M_s =$

s/i	1	2	$\cdots$	n	1	2	$\cdots$	n
1	1	2	$\cdots$	n	1	0	$\cdots$	0
2	1	2	$\cdots$	n	0	1	$\cdots$	0
$\vdots$	1	2	$\cdots$	n	0	0	$\cdots$	0
n	1	2	$\cdots$	n	0	0	$\cdots$	1

(a)



Figure 2:

$f[\lambda_1(s_1, i_1)] = \lambda_2[f(s_1), g(i_1)]$, where $s_j \in S_j$ and $i_j \in I_j$, $j \in \{1, 2\}$. The triple (f, g, h) is an *isomorphism* between M_1 and M_2. An isomorphism just renames the states, the inputs and the outputs. From a purely input/output point of view, g as well as h (or h^{-1}) are combinatory circuits and M_1 performs similar as the serial decomposition (see below) $h^{-1}M_2g$ of the machines g, M_2 and h^1.

The *serial connection* of the two machines $M_1 = (S_1, I_1, O_1, \delta_1, \lambda_1)$ and $M_2 = (S_2, I_2, O_2, \delta_2, \lambda_2)$ for which $O_1 = I_2$ is the machine [13, p. 42]

$$M = M_1 \to M_2 = (S_1 \times S_2, I_1, O_2, \delta, \lambda)$$

where $\delta[(s_1, s_2), i] = (\delta_1(s_1, i), \delta_2[s_2, \lambda(s_1, i)])$ and $\lambda[(s_1, s_2), i] = \lambda_2[s_2, \lambda_1(s, i)]$.

The *parallel connection* of the two machines $M_1 = (S_1, I_1, O_1, \delta_1, \lambda_1)$ and $M_2 = (S_2, I_2, O_2, \delta_2, \lambda_2)$ is the machine [13, p. 48]

$$M = M_1 \| M_2 = (S_1 \times S_2, I_1 \times I_2, O_1 \times O_2, \delta, \lambda)$$

where $\delta[(s_1, s_2), (i_1, i_2)] = (\delta_1(s_1, i_1), \delta_2(s_2, i_2))$ and $\lambda[(s_1, s_2), (i_1, i_2)] = (\lambda_1(s_1, i_1), \lambda_2(s_2, i_2))$.

The logical structure of the initial-state identification problem can be defined as follows. Let us call a proposition concerning the initial state of the machine *experimentally decidable* if there is an experiment E which determines the truth value of that proposition. This can be done by performing E, i.e., by the input of a sequence of input symbols $i_1, i_2, i_3, \ldots, i_n$ associated with E, and by observing the output sequence

$$\lambda_E(s) = \lambda(s, i_1), \lambda(\delta(q, a_1), a_2), \ldots, \lambda(\underbrace{\delta(\cdots\delta(q, a_1)\cdots, a_{n-1})}_{n-1 \text{ times}}, a_n).$$

The most general form of a prediction concerning the initial state s of the machine is that the initial state s is contained in a subset P of the state set S. Therefore, we



Figure 3:

may identify propositions concerning the initial state with subsets of S. A subset P of S is then identified with the proposition that the initial state is contained in P.

Let E be an experiment (a preset or adaptive one), and let $\lambda_E(s)$ denote the obtained output of an initial state s. λ_E defines a mapping of S to the set of output sequences O^*. We define an equivalence relation on the state set S by

$$s \stackrel{E}{\equiv} t \text{ iff } \lambda_E(s) = \lambda_E(t)$$

for any $s, t \in S$. We denote the partition of S corresponding to $\stackrel{E}{\equiv}$ by $S/\stackrel{E}{\equiv}$. Obviously, the propositions decidable by the experiment E are the elements of the Boolean algebra generated by $S/\stackrel{E}{\equiv}$, denoted by B_E.

Let $\mathcal{R}$ be the set of all Boolean algebras B_E. We call the partition logic $(S, \mathcal{R})$ an *automaton propositional calculus.*

Example 5.2 *In what follows, we explicitly construct the Mealy automaton M_s introduced before. Input/output experiments can be performed by the input of one symbol i (in this example, more inputs yield no finer partitions). Let us assume that one input $i = 5$. This experiment is able to distinguish between state $s = 5$ and all the other states; hence it induces a partition (suppose $n > 5$)*

$$v(5) = \{\{5\}, \{1, 2, 3, 4, 6, ..., n\}\}.$$

After this experiment, information about the initial state is lost (irreversible model). Now consider the partitions $v(i)$ of all possible experiments with one input x (all of them non-co-measurable). Every one of them generates a Boolean algebra of events with two atoms; e.g., $v(5)$ generates a two-element Boolean algebra 2^2 whose Hasse diagram is drawn in Fig. 3.

The automaton propositional calculus *and the associated* partition logic *is the set of all partitions*

$$P = \{v(i) \mid i \in I\}.$$

Lattice theoretically, this amounts to a pasting [19] *of all the $v(i)$'s. In the specific example, the pasting is just the horizontal sum—only the least and greatest elements $\emptyset$ and of each 2^2 is identified—and one obtains a Chinese lantern lattice MO_n.*

References

[1] Aerts, A., and Daubechies, I. (1978). Physical justification for using tensor product to describe quantum systems as one joint system *Helvetica Physica Acta,* **51**, 661–675.

[2] G. Boole, (1854). *An Investigation of the Laws of Thought.* Macmillan, reprinted by Dover Press, New York, 1967.

[3] Busch, P., Lahti, P.J., and Mittelstaedt, P. (1991) *The Quantum Theory of Measurement,* Lecture Notes in Physics, Springer–Verlag, Berlin, Heidelberg, New York, London, Budapest.

[4] Dvurečenskij, A. (1995). Tensor product of difference spaces *Transactions of the American Mathematical Society*, **347**, 1043–1057.

[5] Dvurečenskij, A., and Pulmannová, S., (1994). Tensor product of D-posets and D-test spaces. *Reports on Mathematical Physics*, **34**, 4251–275.

[6] Dvurečenskij, A., Pulmannová, S., and Svozil, K. (1995). Partition logics, orthoalgebras and automata. *Helvetica Physica Acta*, **68**, 407–428.

[7] Foulis, D.J. (1989). Coupled physical systems. Foundations of Physics. **19**, 905–922.

[8] Foulis, D.J., and Bennett, M.K. (1993). Tensor products of orthoalgebras, *Order* **10**, 271–282.

[9] Foulis, D.J., Greechie,R.J., and Rüttimann, G.T. (1992). Filters and supports in orthoalgebras, *International Journal of Theoretical Physics,* **31**, 787–807.

[10] Foulis, D., and Pták, P. (1995). On the tensor product of a Boolean algebra and an orthoalgebra, *Czechoslovak Mathematical Journal,* **45 (120)**, 117–126.

[11] Foulis, D., and Randall, C. (1981). Empirical logic and tensor products, In:*Interpretations and Foundations of Quantum Theories,* A. Neumann ed., Wissenschaftsverlag, Bibliographisches Institut, Mannheim, pp. 9–20.

[12] Golfin, A. (1987). *Representations and products of Lattices.* Ph.D. Thesis, Univ. of Massachusetts, Amherst.

[13] Hartmanis, J., and Stearns, R.E. (1966). Algebraic Structure Theory of Sequential Machines. Prentice Hall, Englewood Cliffs, New Jersey.

[14] Hopcroft, J.E., and Ullman, J.D. (1979). *Introduction to Automata Theory, Languages, and Computation.* Addison-Wesley, Reading, Massachussetts.

[15] Kläy, M., Randall, C., and Foulis, D. (1987). Tensor products and probability weights, *International Journal of Theoretical Physics,* **26**, 199–219.

[16] Lock, R. (1990). The tensor product of generalized sample spaces which admit a unital set of dispersion–free weights, *Foundations of Physics,* **20**, 477–498.

[17] Matolcsi, T. (1975). Tensor product of Hilbert lattices and free orthodistributive product of orthomodular lattices, *Acta Scientarum Mathematicarum (Szeged)*, **37**, 263–272.

[18] Moore, E.F. (1956). Gedanken-experiments on sequential machines. In: C. E. Shannon and J. McCarthy, editor, *Automata Studies.* Princeton University Press, Princeton.

[19] Navara, M., and Rogalewicz, V. (1991). The pasting constructions for orthomodular posets. *Mathematische Nachrichten*, **154**, 157–168.

[20] Pták, P., and S. Pulmannová, S. (1991). *Orthomodular Structures as Quantum Logics,* Kluwer, Dordrecht, Boston, London.

[21] Randall, C., and Foulis, D. (1981). Empirical statistics and tensor products. In: *Interpretations and Foundations of Quantum Theory.* H. Neumann ed. vol 5, Wissenschaftsverlag, Bibliographisches Institut, Mannheim, pp. 21–28.

[22] Schaller, M., and Svozil, K. (1994). Partition logics of automata. *Il Nuovo Cimento*, **109B**, 167–176.

[23] Schaller, M., and Svozil, M. (1995). Automaton partition logic versus quantum logic. *International Journal of Theoretical Physics*, **34** (8), 1741–1750.

[24] Schaller,M., and Svozil, K. (1996). Automaton logic. *International Journal of Theoretical Physics*, **35** (to appear).

[25] Svozil, K. (1993). *Randomness and Undecidability in Physics.* World Scientific, Singapore.

[26] Wilce, A. (1990). Tensor product of frame manuals. *International Journal of Theoretical Physics,* **29**, 805–814.

[27] Zecca, A. (1978). On the coupling of quantum logics, *Journal of Mathematical Physics,* **19**, 1482–1485.

This preprint series was founded in 1996. Its purpose is to present manuscripts of submitted or unpublished papers and reports of fellows of the Mathematical Institute of the Slovak Academy of Sciences in Bratislava and its Košice branch. The authors are fully responsible for the content of the preprints. Postal addresses:

Mathematical Institute
Slovak Academy of Sciences
Štefánikova 49
SK–814 73 Bratislava
SLOVAKIA

Tel./Fax (+421 7) 5249 7316

Mathematical Institute
Slovak Academy of Sciences
Grešákova 6
SK–040 01 Košice
SLOVAKIA

Tel./Fax (+421 95) 6228291

PostScript files of preprints are available at WWW address:
`http://www.mat.savba.sk/preprints`

Latest preprints of the series:

14/1996 *Anatolij Dvurečenskij, Karl Svozil*: Product of partition logics, orthoalgebras and automata

13/1996 *Robin L. Hudson, Sylvia Pulmannová*: Chaotic expansion of elements of the universal enveloping algebra of a Lie algebra associated with a quantum stochastic calculus

12/1996 *Sylvia Pulmannová*: Congruences in partial abelian semigroups

11/1996 *Karol Nemoga, Štefan Schwarz*: An explicit description of the set of all normal bases generators of a finite field

10/1996 *Ján Jakubík*: Subdirect product decompositions of MV-algebras

9/1996 *Ivan Korec*: Reversibility in generalized Pascal triangles and binary reversibility in one-dimensional cellular automata

8/1996 *Jozef Bobok, Milan Kuchta*: X-minimal patterns and generalization of Sharkovskiĭ's theorem

7/1996 *Jarmila Hedlíková, Sylvia Pulmannová*: Generalized difference posets and orthoalgebras

6/1996 *Ján Borsík, Roman Frič*: Pointwise convergence fails to be strict

5/1996 *Miroslav Ploščica*: Affine completions of distributive lattices

4/1996 *Ján Haluška*: Equal temperament and Pythagorean tuning: a geometrical interpretation in the plane

3/1996 *Winfried Just, Peter Vojtáš*: On matrix rapid filters

2/1996 *Anatolij Dvurerečenskij*: Fuzzy set representations of some quantum structures

1/1996 *Miloslav Duchoň, Beloslav Riečan*: On the product of semigroup valued measures

(continued on inside back cover)