Mathematical Institute
Slovak Academy of Sciences, Bratislava

Preprint 11/1999

Peter Eliaš

A hierarchy of thin sets related to the boundedness of trigonometric series

July 26, 1999

PREPRINT SERIES

A HIERARCHY OF THIN SETS RELATED TO THE BOUNDEDNESS OF TRIGONOMETRIC SERIES

PETER ELIAŠ

ABSTRACT. We study the family $\mathcal{B}_0$ of the sets on which some series of the form $\sum_{k\in\mathbb{N}} |\sin \pi n_k x|$ is uniformly bounded. We show that the families $\mathcal{B}_0^c$ of all sets admitting the boundary c form a hierarchy which is incontinuous with respect to the operations of intersection and union.

Let us recall two kinds of thin sets studied in harmonic analysis. A set $X \subseteq [0,1]$ is called an *N-set* (in honour of V. V. Nemytskiĭ) if there exists a trigonometric series absolutely converging on X but not converging absolutely everywhere; it is called an N_0*-set* if there exists a series of the form $\sum_{k\in\mathbb{N}} |\sin \pi n_k x|$ converging on X. J. Arbault [1] showed that the family $\mathcal{N}_0$ of all N_0-sets is a proper subfamily of the family $\mathcal{N}$ of all N-sets, and both families share some common properties.

In the paper [3] we examined several modifications of the definitions of the families $\mathcal{N}$ and $\mathcal{N}_0$, and compared the obtained families one to another. We showed that the family $\mathcal{B}_0$ of all sets on which some series of the form $\sum_{k\in\mathbb{N}} |\sin \pi n_k x|$ is uniformly bounded, differs from previously known "classical" families. In the present paper we continue the study of the family $\mathcal{B}_0$ considering the hierarchy of the families $\mathcal{B}_0^c$ of all sets admitting the boundary c. We show that if $c < d$ then $\mathcal{B}_0^c$ is a proper subfamily of $\mathcal{B}_0^d$, and $\bigcup_{d<c} \mathcal{B}_0^d \subsetneqq \mathcal{B}_0^c \subsetneqq \bigcap_{d>c} \mathcal{B}_0^d$ for every $c > 0$.

For a review of families of trigonometric thin sets, some historical notes, and also for many new results we refer the reader to the paper [2].

We shall deal with the quotient group $\mathbb{T} = \mathbb{R}/\mathbb{Z}$; however, we will not distinguish between the elements of $\mathbb{T}$ and the reals, or between the functions defined on $\mathbb{T}$ and the corresponding periodic functions on $\mathbb{R}$. For a real x, let $[x]$ denote the integer part of x and let $\|x\|$ denote the distance of x to the nearest integer, i. e. $[x] = \max\{k \in \mathbb{Z} : k \le x\}$ and $\|x\| = \min\{|x-k| : k \in \mathbb{Z}\}$. Let us note that $\|x\| - \|y\| \le \|x+y\| \le \|x\| + \|y\|$ and $\|-x\| = \|x\|$ for all $x, y \in \mathbb{T}$. The space $\mathbb{T}$ equipped with the metric $\varrho(x,y) = \|x-y\|$ is a compact topological group.

In an accordance with [3], we define B_0-sets as follows.

Definition 1. A set $X \subseteq \mathbb{T}$ is a B_0*-set* if there exists an increasing sequence $\{n_k\}_{k\in\mathbb{N}}$ of natural numbers such that the series $\sum_{k\in\mathbb{N}} \|n_k x\|$ is uniformly bounded on X, i. e. there exists a real c such that $\sum_{k\in\mathbb{N}} \|n_k x\| \le c$ for all $x \in X$. The family of all B_0-sets is denoted by $\mathcal{B}_0$.

1991 *Mathematics Subject Classification.* Primary 43A46; Secondary 42A28.

Key words and phrases. Trigonometric thin sets, N_0-sets, B_0-sets, uniform boundedness.

The work on this paper was supported by grant 2/4034/97 of Slovak Grant Agency VEGA. The research was partly done when the author was visiting the Mathematical Institute of the University in Bonn with financial support by the Graduiertenkolleg.

Since $2\|x\| \le |\sin \pi x| \le \pi \|x\|$ for all $x \in \mathbb{T}$, it is clear that the previous definition remains equivalent if we replace $\sum_{k\in\mathbb{N}} \|n_k x\|$ by $\sum_{k\in\mathbb{N}} |\sin \pi n_k x|$.

Proposition 2. *A set $X \subseteq \mathbb{T}$ is a B_0-set if and only if there exist an increasing sequence $\{n_k\}_{k\in\mathbb{N}}$ of natural numbers and a sequence $\{a_k\}_{k\in\mathbb{N}}$ of elements of $\mathbb{T}$ such that the series $\sum_{k\in\mathbb{N}} \|n_k x + a_k\|$ is uniformly bounded on X.*

Proof. It follows immediately from Lemma 3 below. □

Lemma 3. *Let $\{n_k\}_{k\in\mathbb{N}}$ be an increasing sequence of natural numbers and let $\{a_k\}_{k\in\mathbb{N}}$ be a sequence of elements of $\mathbb{T}$. Then for every $\varepsilon > 0$ there exists an increasing sequence $\{m_j\}_{j\in\mathbb{N}}$ of natural numbers such that $\sum_{j\in\mathbb{N}} \|m_j x\| \le \sum_{k\in\mathbb{N}} \|n_k x + a_k\| + \varepsilon$ for every $x \in \mathbb{T}$.*

Proof. We use a classical argument. Since $\mathbb{T}$ is compact, there exists an increasing function $h : \mathbb{N} \to \mathbb{N}$ such that the sequence $\left\{n_{h(2j+1)} - n_{h(2j)}\right\}_{j\in\mathbb{N}}$ is increasing and $\sum_{j\in\mathbb{N}} \left\|a_{h(2j+1)} - a_{h(2j)}\right\| \le \varepsilon$. It suffices to put $m_j = n_{h(2j+1)} - n_{h(2j)}$. □

However, we do not know the answer for the following question.

Problem 4. Let $\{n_k\}_{k\in\mathbb{N}}$ be an increasing sequence of natural numbers and let $\{a_k\}_{k\in\mathbb{N}}$ be a sequence of elements of $\mathbb{T}$. Does there exist a sequence $\{m_j\}_{j\in\mathbb{N}}$ such that $\sum_{j\in\mathbb{N}} \|m_j x\| \le \sum_{k\in\mathbb{N}} \|n_k x + a_k\|$ for all $x \in \mathbb{T}$?

Let us now define the families $\mathcal{B}_0^c$. Here the use of the series $\sum_{k\in\mathbb{N}} \|n_k x\|$ and $\sum_{k\in\mathbb{N}} |\sin \pi n_k x|$ is not equivalent; for simplicity we will consider the first one.

Definition 5. Let c be a positive real. A set $X \subseteq \mathbb{T}$ is called a *B_0^c-set* if there exists an increasing sequence $\{n_k\}_{k\in\mathbb{N}}$ of natural numbers such that $\sum_{k\in\mathbb{N}} \|n_k x\| \le c$ for all $x \in X$. The family of all B_0^c-sets is denoted by $\mathcal{B}_0^c$.

We show that in the previous definition, '$\le$' can be replaced by '$<$'.

Lemma 6. *Let $c > 0$, $X \subseteq \mathbb{T}$, and let $\{n_k\}_{k\in\mathbb{N}}$ be such that $\sum_{k\in\mathbb{N}} \|n_k x\| \le c$ for all $x \in X$. Then there exists $j \in \mathbb{N}$ such that $\sum_{k>j} \|n_k x\| < c$ for all $x \in X$.*

Proof. Suppose for a contradiction that for every j there exists $x_j \in X$ such that $\sum_{k>j} \|n_k x_j\| \ge c$. Thus $\|n_k x_j\| = 0$ for all $k \le j$ and $\left\|n_{k_j} x_j\right\| > 0$ for some $k_j > j$. Hence for every j and k such that $k \ge k_j$ we have $\left\|n_{k_j} x_j\right\| > 0$ and $\left\|n_{k_j} x_k\right\| = 0$, and thus $x_j \ne x_k$. It follows that the set $\{x_j : j \in \mathbb{N}\}$ is infinite. However, this is impossible since for every k there are only finitely many $x \in \mathbb{T}$ such that $\|n_k x\| = 0$. □

Let us recall that a set $X \subseteq \mathbb{T}$ is called a *Dirichlet set* (a *D-set*) if there exists an increasing sequence $\{n_k\}_{k\in\mathbb{N}}$ of natural numbers such that the sequence $\{\|n_k x\|\}_{k\in\mathbb{N}}$ is uniformly converging to 0 on X. The family of all D-sets is denoted by $\mathcal{D}$. From the definition we can immediately see that $\bigcup_{c>0} \mathcal{B}_0^c = \mathcal{B}_0$ and $\bigcap_{c>0} \mathcal{B}_0^c = \mathcal{D}$.

We do not know whether an analogue of Proposition 2 holds true in the case of B_0^c-sets.

Problem 7. Let $c > 0$, let $\{n_k\}_{k\in\mathbb{N}}$ be an increasing sequence of natural numbers and let $\{a_k\}_{k\in\mathbb{N}}$ be a sequence of elements of $\mathbb{T}$. Does there exist a sequence $\{m_j\}_{j\in\mathbb{N}}$ such that for all $x \in \mathbb{T}$, $\sum_{j\in\mathbb{N}} \|m_j x\| \le c$ whenever $\sum_{k\in\mathbb{N}} \|n_k x + a_k\| \le c$?

Fix a positive real c. Let us denote by $\mathcal{B}_{/}^{*\rfloor}$ the family of all sets $X \subseteq \mathbb{T}$ for which there exist sequences $\{n_k\}_{k\in\mathbb{N}}$ and $\{a_k\}_{k\in\mathbb{N}}$ such that $\sum_{k\in\mathbb{N}} \|n_k x + a_k\| \leq c$ for all $x \in X$. Clearly $\mathcal{B}_{/}^{\rfloor} \subseteq \mathcal{B}_{/}^{*\rfloor}$, and Problem 7 asks whether the equality holds true. From Lemma 3 it follows that $\mathcal{B}_{/}^{*\rfloor} \subseteq \bigcap_{\lceil>\rfloor} \mathcal{B}_{/}^{\lceil}$. Thus for every $c > 0$ we have

$$\bigcup_{d<c} \mathcal{B}_{/}^{\lceil} = \bigcup_{\lceil<\rfloor} \mathcal{B}_{/}^{*\lceil} \subseteq \mathcal{B}_{/}^{\rfloor} \subseteq \mathcal{B}_{/}^{*\rfloor} \subseteq \bigcap_{\lceil>\rfloor} \mathcal{B}_{/}^{\lceil} = \bigcap_{\lceil>\rfloor} \mathcal{B}_{/}^{*\lceil}.$$

In Propositions 10 and 11 we will show that the first and the last inclusions are proper.

Let $\{m_j\}_{j\in\mathbb{N}}$ be an increasing sequence of natural numbers and let $\{a_j\}_{j\in\mathbb{N}}$ be a sequence of elements of $\mathbb{T}$. For every positive integer n, let $\alpha(n), \beta(n) \in \mathbb{N}$ be such that $n = 2^{\alpha(n)}\beta(n)$ and $\beta(n)$ is odd.

Lemma 8. *Let $p, r \in \mathbb{N}$, $p \geq 1$. Assume that the set $\{\alpha(m_j) : j \in \mathbb{N}\}$ is infinite. There exist $j, q \in \mathbb{N}$ and $s \in \{-1, 1\}$ such that $q \geq r$ and $\left\|sm_j 2^{-q2^p} + a_j\right\| \geq 2^{-2^p}$.*

Proof. Put $J = \{j \in \mathbb{N} : \|a_j\| \geq 2^{-2^p}\}$. Clearly either $\{\alpha(m_j) : j \in J\}$ or $\{\alpha(m_j) : j \notin J\}$ is infinite.

In the first case find $j \in J$ such that $\alpha(m_j) \geq r2^p$, and put $q = r$ and $s = 1$. Then $sm_j 2^{-q2^p}$ is an integer and thus $\left\|sm_j 2^{-q2^p} + a_j\right\| = \|a_j\| \geq 2^{-2^p}$.

In the latter case find $j \notin J$ such that $\alpha(m_j) \geq (r-1)2^p$ and put $q = [\alpha(m_j)2^{-p}] + 1$. Then clearly $q \geq r$. We also have $0 < q2^p - \alpha(m_j) \leq 2^p$ and thus $\left\|m_j 2^{-q2^p}\right\| = \left\|\beta(m_j)2^{\alpha(m_j)-q2^p}\right\| \geq 2^{-2^p}$. Since $\|a_j\| \leq 2^{-2^p} \leq 2^{-2}$, we can conclude that either $\left\|m_j 2^{-q2^p} + a_j\right\| \geq 2^{-2^p}$ or $\left\|m_j 2^{-q2^p} - a_j\right\| \geq 2^{-2^p}$. $\square$

Lemma 9. *Let θ be a real such that $0 < \theta \leq 1/2$, let $\{p_k\}_{k\in\mathbb{N}}$ be a sequence of positive integers, and let $\{\varepsilon_k\}_{k\in\mathbb{N}}$ be a sequence of positive reals. Assume that $\lim_{j\to\infty} \alpha(m_j) = \infty$. There exists $x \in \mathbb{T}$ such that*

1. *$\|x\| \leq \theta$;*
2. *there exists an increasing sequence $\{n_k\}_{k\in\mathbb{N}}$ of natural numbers such that $\left\|2^{2^{n_k}} x\right\| \leq 2^{-2^{p_k}} + 2^{-2^{n_k}}$ for all $k \in \mathbb{N}$, and $\left\|2^{2^n} x\right\| \leq 2^{-2^n}$ for all $n \notin \{n_k : k \in \mathbb{N}\}$;*
3. *there exists an increasing sequence $\{j_k\}_{k\in\mathbb{N}}$ of natural numbers such that $\|m_{j_k} x + a_{j_k}\| \geq 2^{-2^{p_k}} - \varepsilon_k$ for all $k \in \mathbb{N}$.*

Proof. We will define sequences $\{n_k\}_{k\in\mathbb{N}}$, $\{j_k\}_{k\in\mathbb{N}}$, $\{q_k\}_{k\in\mathbb{N}}$, and $\{s_k\}_{k\in\mathbb{N}}$ by induction. Fix $k \in \mathbb{N}$ and suppose that n_i, j_i, q_i, s_i are defined for all $i < k$. Find $r \geq 2$ such that

(a) $2^{-r2^{p_k}} \leq 2^{-k-1}\theta$;
(b) $r2^{p_k} > 2^{n_{k-1}+1}$ if $k \geq 1$;
(c) $2^{-r2^{p_k}} \leq 2^{i-k} 2^{-2^{n_i+1}}$ for all $i < k$;
(d) $2^{-r2^{p_k}} \leq 2^{i-k}\varepsilon_i / m_{j_i}$ for all $i < k$.

Let $l \in \mathbb{N}$ be such that $l > j_{k-1}$ if $k \geq 1$, and $\alpha(m_j) \geq q_{k-1}2^{p_{k-1}}$ for all $j \geq l$. By Lemma 8 there exist $j_k \geq l$, $q_k \geq r$, and $s_k \in \{-1, 1\}$ such that

(e) $\left\|m_{j_k} s_k 2^{-q_k 2^{-p_k}} + a_{j_k}\right\| \geq 2^{-2^{p_k}}$.

Put $n_k = \max\{i : 2^i < q_k 2^{p_k}\}$.

Let $x = \sum_{k\in\mathbb{N}} s_k 2^{-q_k 2^{p_k}}$. By (a), $|x| \leq \sum_{k\in\mathbb{N}} 2^{-q_k 2^{p_k}} \leq \sum_{k\in\mathbb{N}} 2^{-k-1}\theta = \theta$, and thus (1) holds.

To show (2), fix $k \in \mathbb{N}$. If $k \geq 1$ then by (b), $q_k 2^{p_k} > 2^{n_{k-1}+1}$. Thus $n_k > n_{k-1}$ and hence $q_i 2^{p_i} < 2^{n_k}$ for all $i < k$. Since $q_k \geq 2$, we have $2^{n_k+1} \geq q_k 2^{p_k} \geq 2^{p_k+1}$ and thus $n_k \geq p_k$. Hence 2^{n_k} is a multiple of 2^{p_k} and we have $q_k 2^{p_k} \geq 2^{n_k} + 2^{p_k}$. Using (c) we obtain that

$$\begin{aligned} \sum_{i \geq k} 2^{-q_i 2^{p_i}} &\leq 2^{-q_k 2^{p_k}} + \sum_{i>k} 2^{k-i} 2^{-2^{n_k+1}} \\ &\leq 2^{-2^{n_k} - 2^{p_k}} + 2^{-2^{n_k+1}} = 2^{-2^{n_k}} \left(2^{-2^{p_k}} + 2^{-2^{n_k}}\right). \end{aligned} \tag{f}$$

Hence $\left\|2^{2^{n_k}} x\right\| \leq 2^{2^{n_k}} \sum_{i \geq k} 2^{-q_i 2^{p_i}} \leq 2^{-2^{p_k}} + 2^{-2^{n_k}}$, and thus the first part of (2) holds. To show the latter one, let $n \notin \{n_k : k \in \mathbb{N}\}$ and put $k = \min\{i : n_i > n\}$. Then for all $i < k$ we have $n > n_i$ and thus $q_i 2^{p_i} \leq 2^n$. Hence by (f),

$$\left\|2^{2^n} x\right\| \leq 2^{2^n} \sum_{i \geq k} 2^{-q_i 2^{p_i}} \leq 2^{2^n - 2^{n_k}} \left(2^{-2^{p_k}} + 2^{-2^{n_k}}\right) \leq 2^{2^n - 2^{n+1}} = 2^{-2^n}.$$

To show (3), fix again $k \in \mathbb{N}$. From the choice of l it follows that $j_k > j_{k-1}$ if $k \geq 1$, and $\alpha(m_{j_k}) \geq q_i 2^{p_i}$ for all $i < k$. Hence by (e) and (d),

$$\begin{aligned} \|m_{j_k} x + a_{j_k}\| &\geq \left\|m_{j_k} s_k 2^{-q_k 2^{p_k}} + a_{j_k}\right\| - m_{j_k} \sum_{i>k} 2^{-q_i 2^{p_i}} \\ &\geq 2^{-2^{p_k}} - m_{j_k} \sum_{i>k} 2^{k-i} \varepsilon_k / m_{j_k} = 2^{-2^{p_k}} - \varepsilon_k. \end{aligned}$$

□

For $n \in \mathbb{N}$, denote $\phi(n) = \sum_{k \geq n} 2^{-2^k}$, $\psi(0) = 1$, and $\psi(n) = 2^{-2^n} \sum_{k<n} 2^{2^k}$ if $n \geq 1$. It can be easily shown that both $\{\phi(n)\}_{n \in \mathbb{N}}$ and $\{\psi(n)\}_{n \in \mathbb{N}}$ are decreasing and converging to 0.

Proposition 10. *For every $c > 0$ there exists $X \subseteq \mathbb{T}$ such that*

1. *there exists an increasing sequence $\{n_k\}_{k \in \mathbb{N}}$ of natural numbers such that $\sum_{k \in \mathbb{N}} \|n_k x\| \leq c$ for all $x \in X$;*
2. *for every increasing sequence $\{m_j\}_{j \in \mathbb{N}}$ of natural numbers, every sequence $\{a_j\}_{j \in \mathbb{N}}$ of elements of $\mathbb{T}$, and for every $\eta > 0$ there exists $x \in X$ such that $\sum_{j \in \mathbb{N}} \|m_j x + a_j\| > c - \eta$.*

Proof. Put $X = \left\{x \in \mathbb{T} : \sum_{\ltimes \in \mathbb{N}} : \left\|\nvDash^{\nvDash^{\ltimes}} \curvearrowleft\right\| \leq \right\}$. Clearly X satisfies the condition (1).

To show (2), fix $\eta > 0$ and sequences $\{m_j\}_{j \in \mathbb{N}}$, $\{a_j\}_{j \in \mathbb{N}}$. We will find $x \in X$ such that $\sum_{j \in \mathbb{N}} \|m_j x + a_j\| > c - \eta$. We will consider two cases.

(a) If $\lim_{j \to \infty} \alpha(m_j) = \infty$ then we can find $N \in \mathbb{N}$, a sequence $\{p_k\}_{k \in \mathbb{N}}$ of positive integers and a sequence $\{\varepsilon_k\}_{k \in \mathbb{N}}$ of positive reals such that $\sum_{k \in \mathbb{N}} 2^{-2^{p_k}} - \sum_{k \in \mathbb{N}} \varepsilon_k > c - \eta$ and $\sum_{k \in \mathbb{N}} 2^{-2^{p_k}} + \phi(N) + \psi(N) \leq c$. Put $\theta = 2^{-2^N}$. There exists $x \in \mathbb{T}$ satisfying the conditions (1)–(3) of Lemma 9. We have

$$\sum_{n \in \mathbb{N}} \left\|2^{2^n} x\right\| = \sum_{n<N} \left\|2^{2^n} x\right\| + \sum_{n \geq N} \left\|2^{2^n} x\right\| \leq \sum_{n<N} 2^{2^n} \theta + \sum_{k \in \mathbb{N}} 2^{-2^{p_k}} + \sum_{n \geq N} 2^{-2^n} \leq c$$

and

$$\sum_{j \in \mathbb{N}} \|m_j x + a_j\| \geq \sum_{k \in \mathbb{N}} \|m_{j_k} x + a_{j_k}\| \geq \sum_{k \in \mathbb{N}} \left(2^{-2^{p_k}} - \varepsilon_k\right) > c - \eta.$$

(b) If the sequence $\{\alpha(m_j)\}_{j\in\mathbb{N}}$ does not tend to infinity then we will find $x \in X$ such that the sequence $\{\|m_j x + a_j\|\}_{j\in\mathbb{N}}$ does not tend to 0. We can suppose that $\lim_{j\to\infty}\|a_j\| = 0$ (otherwise take $x = 0$). There exist $M \in \mathbb{N}$ and an increasing sequence $\{j_k\}_{k\in\mathbb{N}}$ of natural numbers such that $\alpha(m_{j_k}) = M$ for all k. Find $N \in \mathbb{N}$ such that $2^N > M$ and $\psi(N) \le c$. For $x = 2^{-2^N}$ we get

$$\sum_{n\in\mathbb{N}} \left\|2^{2^n} x\right\| = \sum_{n<N} 2^{2^n - 2^N} = \psi(N) \le c.$$

Since $\|m_{j_k} x + a_{j_k}\| \ge \|m_{j_k} x\| - \|a_{j_k}\| \ge 2^{M-2^N} - \|a_{j_k}\|$ for all $k \in \mathbb{N}$, the sequence $\{\|m_j x + a_j\|\}_{j\in\mathbb{N}}$ does not converge to 0. $\square$

Proposition 11. *For every $c > 0$ there exists $X \subseteq \mathbb{T}$ such that*

1. *for every $\eta > 0$ there exists an increasing sequence of natural numbers $\{n_k\}_{k\in\mathbb{N}}$ such that $\sum_{k\in\mathbb{N}}\|n_k x\| < c + \eta$ for all $x \in X$;*
2. *for every increasing sequence $\{m_j\}_{j\in\mathbb{N}}$ of natural numbers and for every sequence $\{a_j\}_{j\in\mathbb{N}}$ of elements of $\mathbb{T}$ there exists $x \in X$ such that $\sum_{j\in\mathbb{N}}\|m_j x + a_j\| \ge c$.*

Proof. Put

$$X = \left\{x \in \mathbb{T} : \sum_{n\ge N}\left\|2^{2^n} x\right\| \le c + \phi(N) + \psi(N) \text{ for all } N \in \mathbb{N}\right\}.$$

Clearly X satisfies the condition (1).

To show (2), fix sequences $\{m_j\}_{j\in\mathbb{N}}$, $\{a_j\}_{j\in\mathbb{N}}$. We will find $x \in X$ such that $\sum_{j\in\mathbb{N}}\|m_j x + a_j\| \ge c$. Again let us discuss two cases.

(a) Assume that $\lim_{j\to\infty}\alpha(m_j) = \infty$. Let us choose a sequence $\{p_k\}_{k\in\mathbb{N}}$ of positive integers such that $\sum_{k\in\mathbb{N}} 2^{-2^{p_k}} = c$. Find $t \in \mathbb{N}$ such that $\left\|m_0 2^{-2^t} + a_0\right\| > 0$ and put $\theta = \left\|m_0 2^{-2^t} + a_0\right\|/2$. Fix a sequence $\{\varepsilon_k\}_{k\in\mathbb{N}}$ of positive reals such that $\sum_{k\in\mathbb{N}}\varepsilon_k = \theta$ and find $l \in \mathbb{N}$ such that $\alpha(m_j) \ge 2^t$ for all $j \ge l$. By Lemma 9 there exists $x' \in \mathbb{T}$ such that

(1') $\|x'\| \le \theta/m_0$;

(2') there exists an increasing sequence $\{n_k\}_{k\in\mathbb{N}}$ of natural numbers such that $\left\|2^{2^{n_k}} x'\right\| \le 2^{-2^{p_k}} + 2^{-2^{n_k}}$ for all $k \in \mathbb{N}$, and $\left\|2^{2^n} x'\right\| \le 2^{-2^n}$ for all $n \notin \{n_k : k \in \mathbb{N}\}$;

(3') there exists an increasing sequence $\{j_k\}_{k\in\mathbb{N}}$ of natural numbers such that $j_k \ge l$ and $\|m_{j_k} x' + a_{j_k}\| \ge 2^{-2^{p_k}} - \varepsilon_k$ for all $k \in \mathbb{N}$.

Put $x = x' + 2^{-2^t}$. Then for every $N \in \mathbb{N}$ we have

$$\begin{aligned}\sum_{n\ge N}\left\|2^{2^n} x\right\| &\le \sum_{n\ge N}\left\|2^{2^n} x'\right\| + \sum_{n\ge N}\left\|2^{2^n - 2^t}\right\| \\ &\le \sum_{k\in\mathbb{N}} 2^{-2^{p_k}} + \sum_{n\ge N} 2^{-2^n} + \sum_{N\le n<t} 2^{2^n - 2^t} \le c + \phi(N) + \psi(N),\end{aligned}$$

since the last sum equals 0 if $t \leq N$ and is not greater than $\psi(t) < \psi(N)$ if $t > N$. Moreover,

$$\begin{aligned}\sum_{j \in \mathbb{N}} \|m_j x + a_j\| &\geq \|m_0 x + a_0\| + \sum_{k \in \mathbb{N}} \|m_{j_k} x + a_{j_k}\| \\ &\geq \left\| m_0 2^{-2^t} + a_0 \right\| - \|m_0 x'\| + \sum_{k \in \mathbb{N}} \|m_{j_k} x' + a_{j_k}\| \\ &\geq \theta + \sum_{k \in \mathbb{N}} 2^{-2^{p_k}} - \sum_{k \in \mathbb{N}} \varepsilon_k = c.\end{aligned}$$

(b) If the sequence $\{\alpha(m_j)\}_{j \in \mathbb{N}}$ does not tend to infinity then similarly as in the proof of Proposition 10 there exists $x \in \mathbb{T}$ such that $\sum_{n \in \mathbb{N}} \left\| 2^{2^n} x \right\| \leq c$ and the sequence $\{\|m_j x + a_j\|\}_{j \in \mathbb{N}}$ does not tend to 0. $\square$

The following theorem summarizes our results concerning the hierarchy of the families $\mathcal{B}_{/}^{\rfloor}$ and $\mathcal{B}_{/}^{*\rfloor}$.

Theorem 12. *Let c be a positive real. Then*

$$\bigcup_{d<c} \mathcal{B}_{/}^{\lceil} = \bigcup_{\lceil < \rfloor} \mathcal{B}_{/}^{*\lceil} \subsetneqq \mathcal{B}_{/}^{\rfloor} \subseteq \mathcal{B}_{/}^{*\rfloor} \subsetneqq \bigcap_{\lceil > \rfloor} \mathcal{B}_{/}^{\lceil} = \bigcap_{\lceil > \rfloor} \mathcal{B}_{/}^{*\lceil}.$$

Corollary 13. *If $c < d$ then $\mathcal{B}_{/}^{\rfloor} \subsetneqq \mathcal{B}_{/}^{\lceil}$ and $\mathcal{B}_{/}^{*\rfloor} \subsetneqq \mathcal{B}_{/}^{*\lceil}$.*

Let us note that the proofs of Propositions 10 and 11 can be easily adopted for showing the following statements.

Proposition 14. *For every $c > 0$ there exists $X \subseteq \mathbb{T}$ such that*

1. *there exists an increasing sequence $\{n_k\}_{k \in \mathbb{N}}$ of natural numbers such that $\sum_{k \in \mathbb{N}} |\sin \pi n_k x| \leq c$ for all $x \in X$;*
2. *for every increasing sequence $\{m_j\}_{j \in \mathbb{N}}$ of natural numbers, every sequence $\{a_j\}_{j \in \mathbb{N}}$ of elements of $\mathbb{T}$, and for every $\eta > 0$ there exists $x \in X$ such that $\sum_{j \in \mathbb{N}} |\sin \pi(m_j x + a_j)| > c - \eta$.*

Proposition 15. *For every $c > 0$ there exists $X \subseteq \mathbb{T}$ such that*

1. *for every $\eta > 0$ there exists an increasing sequence of natural numbers $\{n_k\}_{k \in \mathbb{N}}$ such that $\sum_{k \in \mathbb{N}} |\sin \pi n_k x| < c + \eta$ for all $x \in X$;*
2. *for every increasing sequence $\{m_j\}_{j \in \mathbb{N}}$ of natural numbers and for every sequence $\{a_j\}_{j \in \mathbb{N}}$ of elements of $\mathbb{T}$ there exists $x \in X$ such that $\sum_{j \in \mathbb{N}} |\sin \pi(m_j x + a_j)| \geq c$.*

Hence it is possible to prove results analogous to Theorem 12 and Corollary 13, concerning the families of sets obtained by replacing the expression $\sum_{k \in \mathbb{N}} \|n_k x\|$ in Definition 5 by $\sum_{k \in \mathbb{N}} |\sin \pi n_k x|$.

References

[1] J. Arbault, *Sur l'ensemble de convergence absolue d'une série trigonométrique*, Bull. Soc. Math. France **80** (1952), 253–317.
[2] L. Bukovský, N. N. Kholshchevnikova and M. Repický, *Thin sets of harmonic analysis and infinite combinatorics*, Real Anal. Exchange **20** (1994/95), 454–509.
[3] P. Eliaš, *A classification of trigonometrical thin sets and their interrelations*, Proc. Amer. Math. Soc. **125** (1997), 1111–1121.

Mathematical Institute, Slovak Academy of Sciences, Jesenná 5, 041 54 Košice, Slovakia
E-mail address: elias@kosice.upjs.sk

This preprint series was founded in 1996. Its purpose is to present manuscripts of submitted or unpublished papers and reports of fellows of the Mathematical Institute of the Slovak Academy of Sciences in Bratislava and its Košice branch. The authors are fully responsible for the content of the preprints. Postal addresses:

Mathematical Institute
Slovak Academy of Sciences
Štefánikova 49
SK–814 73 Bratislava
SLOVAKIA

Tel./Fax (+421 7) 5249 7316

Mathematical Institute
Slovak Academy of Sciences
Grešákova 6
SK–040 01 Košice
SLOVAKIA

Tel./Fax (+421 95) 6228291

PostScript files of preprints are available at WWW address:
`http://www.mat.savba.sk/preprints`

Latest preprints of the series:

11/1999 *Peter Eliaš*: A hierarchy of thin sets related to the boundedness of trigonometric series

10/1999 *Miloslav Duchoň, Peter Maličký*: On Herglotz theorem in partially ordered vector spaces

9/1999 *Ján Jakubík*: Torsion classes of specker lattice ordered groups

8/1999 *Ján Jakubík*: Graph automorphisms and cells of lattices

7/1999 *Miloslav Duchoň, Peter Maličký*: Positive linear maps on trigonometric polynomials

6/1999 *Miloslav Duchoň, Anna Kolesárová*: Fourier transforms of vector-valued measures on a certain compact semigroup

5/1999 *Miroslav Ploščica*: Separation properties in congruence lattices of lattices

4/1999 *Georges Chevalier, Anatolij Dvurečenskij, Karl Svozil*: Piron's and Bell's geometric lemmas and Gleason's theorem

3/1999 *Jarmila Hedlíková*: Relatively orthomodular lattices

2/1999 *Anatolij Dvurečenskij, Pekka Lahti, Kari Ylinen*: Some remarks on the moment problem

1/1999 *Anatolij Dvurečenskij*: Loomis-Sikorski theorem for σ-complete MV-algebras and ℓ-groups

27/1998 *Martin Kochol*: Cubic graphs without a Petersen minor have nowhere-zero 5-flows

26/1998 *Anatolij Dvurečenskij, Ferdinand Chovanec, Eva Rybáriková*: D-homomorphisms and atomic σ-complete Boolean D-posets

25/1998 *Anatolij Dvurečenskij*: Commutative BCK-algebras with product

24/1998 *Anatolij Dvurečenskij, Antonio Di Nola*: Product MV-algebras

23/1998 *Anatolij Dvurečenskij, Maria Gabriella Graziano*: Dedekind complete commutative BCK-algebras

22/1998 *Anatolij Dvurečenskij*: Ideals of commutative BCK-algebras

21/1998 *Anatolij Dvurečenskij, Beloslav Riečan*: Weakly divisible MV-algebras and product

20/1998 *Ivan Korec*: Definability of addition and multiplication from Pascal's triangle modulo n and number-theoretical functions

19/1998 *Roberto Giuntini, Sylvia Pulmannová*: Ideals and congruences in effect algebras and QMV algebras

18/1998 *Ivan Korec*: Definability of exact arithmetical operations from approximate ones

(continued on inside back cover)

17/1998 *Georges Chevalier, Sylvia Pulmannová*: Some ideal lattices in partial abelian monoids

16/1998 *Anatolij Dvurečenskij*: On categorical equivalences of commutative BCK-algebras

15/1998 *Ivan Korec*: Definability of addition and multiplication from some quadratic forms $ax^2 + cy^2$

14/1998 *Beloslav Riečan*: On the L^p space of observables on product MV algebras

13/1998 *Beloslav Riečan*: On the strong law of large numbers on some ordered structures

12/1998 *Ladislav Mišík, jr., Tibor Žáčik*: A simplified formula for calculation of metric dimension of converging sequences

11/1998 *Beloslav Riečan*: On the joint distribution of observables

10/1998 *Beloslav Riečan*: On the product MV algebras

9/1998 *Roman Frič*: A Stone-type duality and its applications to probability

8/1998 *Roman Frič*: Ring of maps: sequential convergence and completion

7/1998 *Roman Frič, Ján Jakubík*: Sequential convergences on Boolean algebras defined by systems of maximal filters

6/1998 *Roman Frič*: Categorical reflections on fields of events and probability

5/1998 *Beloslav Riečan*: On the probability theory and fuzzy sets

4/1998 *Ivan Korec*: Definability of arithmetical operations from binary quadratic forms

3/1998 *Beloslav Riečan*: On the conditional expectation of observables in MV algebras of fuzzy sets

2/1998 *Beloslav Riečan*: On the conditional probability on MV algebras

1/1998 *Paula B. Cohen, Robin L. Hudson, K. R. Parthasarathy, S. Pulmannová*: Hall's transformation via quantum stochastic calculus

13/1997 *Ivan Korec*: Real-time generation of primes by a one-dimensional cellular automaton with 9 states

12/1997 *Ivan Korec*: Arithmetical operations strongest with respect to the first order definability

11/1997 *Otokar Grošek, Karol Nemoga, Marcel Zanechal*: Why use bijective S-boxes in GOST-algorithm

10/1997 *Beloslav Riečan*: On the L^p space of observables

9/1997 *Ladislav Mišík Jr., Tibor Žáčik*: A formula for calculation of metric dimension of converging sequences

8/1997 *Anatolij Dvurečenskij, Maria Gabriella Graziano*: Commutative *BCK*-algebras and lattice ordered groups

7/1997 *Anatolij Dvurečenskij, Maria Gabriella Graziano*: On representations of commutative BCK-algebras

6/1997 *Sylvia Pulmannová, Karl Svozil*: Ideals in ortholattices, Bell inequalities and simultaneously definite properties

5/1997 *Camille Debiève, Miloslav Duchoň, Beloslav Riečan*: Moment problem in some ordered vector spaces

4/1997 *Miloslav Duchoň, Beloslav Riečan*: On the Kurzweil–Stieltjes integral in ordered spaces

3/1997 *Anatolij Dvurečenskij, Maria Gabriella Graziano*: Remarks on representations of minimal clans

2/1997 *Ivan Korec*: Elementary definability from Pascal's triangle modulo p and the set of e-th powers

1/1997 *Ján Jakubík*: Lexicographic products of half linearly ordered groups

34/1996 *Emília Halušková*: Direct limits of monounary algebras