Mathematical Institute
Slovak Academy of Sciences, Bratislava

Preprint 3/2002

Ján Jakubík

Generalized cardinal properties of lattices and lattice ordered groups

January 17, 2002

PREPRINT SERIES

CANTOR-BERNSTEIN THEOREM FOR LATTICES

Ján Jakubík, Košice

Abstract. This paper is a continuation of the author's article [6]; the result of [6] is extended to the case when the lattice under consideration need not have the least element.

In the paper [6] a result of Cantor-Bernstein type was proved for lattices which (a) have the least element, (b) are σ-complete, and (c) are infinitely distributive.

In the present paper we modify the method from [6] to obtain a generalization of the mentioned result such that the condition (a) is omitted and the conditions (b), (c) are substantially weakened.

We remark that a theorem of Sikorski [10] (proved independently by Tarski [13], cf. also Sikorski [11]) concerning σ-complete Boolean algebras is a corollary of the result from [6].

1. Preliminaries

We denote by $\mathcal{T}_\sigma^0$ the class of all lattices satisfying the conditions (a), (b) and (c) above.

Let L be a lattice and $x_0 \in L$. An indexed system $(x_i)_{i\in I}$ of elements of L will be called orthogonal over x_0 if (i) $x_i \geqq x_0$ for each $i \in I$, and (ii) $x_{i(1)} \wedge x_{i(2)} = x_0$ whenever $i(1)$ and $i(2)$ are distinct elements of I. The orthogonality under x_0 is defined dually.

Let α be an infinite cardinal. Consider the following conditions for L:

(b'_α) If $x_0 \in L$ and $(x_i)_{i\in I}$ is an indexed system of elements of L which is orthogonal over x_0 and if $\operatorname{card} I \leqq \alpha$, then the join $\bigvee_{i\in I} x_i$ exists in L.

(b''_α) If the assumption of (b'_α) is satisfied and if, moreover, the system $(x_i)_{i\in I}$ is upper bounded in L, then $\bigvee_{i\in I} x_i$ exists in L.

(c'_α) If the assumption of (b'_α) is satisfied and if, moreover, the join $\bigvee_{i\in I} x_i$ exists in L, then for each Selement $y \in L$ the relation

$$y \wedge (\bigvee_{i\in I} x_i) = \bigvee_{i\in I} (y \wedge x_i)$$

is valid in L.

1991 *Mathematics Subject Classification.* 06B05.

Key words and phrases. Lattice, direct product decomposition, Cantor-Bernstein Theorem.

Supported by grant VEGA 2/5125/98

We denote by $(\mathrm{b}'_{\alpha d})$, $(\mathrm{b}''_{\alpha d})$ and $(\mathrm{c}'_{\alpha d})$ the conditions which are dual to (b'_{α}), (b''_{α}) or (c'_{α}), respectively.

Let $\mathcal{T}^1_\alpha$ be the class of all lattices which satisfy the conditions (b''_{α}), (c'_{α}), $(\mathrm{b}''_{\alpha d})$ and $(\mathrm{c}'_{\alpha d})$. Next let $\mathcal{T}^2_\alpha$ be the class of all lattices satisfying (b'_{α}), (c'_{α}), $(\mathrm{b}'_{\alpha d})$ and $(\mathrm{c}'_{\alpha d})$.

We use the notion of internal direct factor with a given central element of a lattice in the same sense as in [6].

The main results of the present paper are Theorem 2.7 and Theorem 3.8. The first one of these theorems says that if $L \in \mathcal{T}^1_\alpha$ and $x^0 \in L$, then the Boolean algebra of all internal direct factors of L with the central element s^0 is α-complete. Theorem 3.8 is a result of Cantor-Bernstein type for lattices belonging to $\mathcal{T}^2_\alpha$, where $\alpha = \aleph_0$; this result is stronger than Theorem 2 of [6].

We substiantially apply the methods from [6].

Some theorems of Cantor-Bernstein type for lattice ordered groups and for MV-algebras were proved in [1]–[5], [7]–[9].

2. Internal direct product decompositions

Let L be a lattice belonging to $\mathcal{T}^1_\alpha$, where α is an infinite cardinal. Further let s^0 be an arbitrary but fixed element of L.

We use the terminology and the notation as in [5]; the reader is assumed to be acquainted with the results of Section 2 of [5].

Let I be a set with $\operatorname{card} I = \alpha$. Assume that for each $i \in I$ we have an internal direct product decomposition

$$L = (s^0)L_i \times L'_i \tag{1}$$

with the central element s^0. We suppose that whenever $i(1)$ and $i(2)$ are distinct elements of I, then

$$L_{i(1)} \cap L_{i(2)} = \{s^0\}. \tag{2}$$

For $x \in L$ and $i \in I$ we denote

$$x_i = x(L_i), \quad x'_i = x(L'_i).$$

Let $x, y \in L$. We put xR_iy if $x(L'_i) = y(L'_i)$. Analogously, we set xR'_iy if $x(L_i) = y(L_i)$. Then R_i and R'_i are permutable congruence relations on L with $R_i \wedge R'_i = R_{\min}$ and $R_i \vee R'_i = R_{\max}$.

For each congruence relation ρ on L and each $x \in L$ we put

$$x_\rho = \{y \in L : x\rho y\}.$$

Then we have

$$L_i = s^0_{R'_i}, \quad L'_i = s^0_{R_i}, \quad \{x(L_i)\} = s^0_{R_i} \cap x_{R'_i}, \quad \{x(L'_i)\} = s^0_{R'_i} \cap x_{R_i}. \tag{3}$$

We shall systematically apply the relations (3).

2.1. Lemma. *Let $x^0 \in L$. Then*

$$x^0_{R_{i(1)}} \cap x^0_{R_{i(2)}} = \{x^0\}$$

whenever $i(1)$ and $i(2)$ are distinct elements of I.

Proof. This is an immediate consequence of (3). □

Let $a, b \in L$, $a \leqq b$. Further let $i \in I$. There exist uniquely determined elements x^i and y^i in L such that

$$(x^i)_i = b_i, \quad (x^i)'_i = a'_i,$$
$$(y^i)_i = a_i, \quad (y^i)'_i = b_i.$$

Then

$$\{x^i\} = a_{R_i} \cap b_{R'_i}, \quad \{y^i\} = a_{R'_i} \cap b_{R_i}. \tag{4}$$

From the definition of x^i and y^i we obtain

$$x^i, y^i \in [a, b] \quad \text{for each } i \in I. \tag{5}$$

2.2. Lemma. *Let $i(1)$ and $i(2)$ be distinct elements of I. Then*

$$x^{i(1)} \wedge x^{i(2)} = a, \quad y^{i(1)} \vee y^{i(2)} = b.$$

Proof. Put $x^{i(1)} \wedge x^{i(2)} = t$. In view of (5), $t \geqq a$. Then $t \in [a, x^{i(1)}]$ and hence according to (4), $t \in a_{R_{i(1)}}$; similarly, $t \in a_{R_{i(2)}}$. Thus 2.1 yields that $t = a$. Therefore $x^{i(1)} \wedge x^{i(2)} = a$. Analogously we obtain $y^{i(1)} \vee y^{i(2)} = b$. □

2.3. Corollary. *Under the notation as above, the indexed system $(x^i)_{i\in I}$ is orthogonal over a, and the indexed system $(y^i)_{i\in I}$ is orthogonal under b.*

Since these systems are bounded, we get

2.4. Corollary. *There exist elements x and y in L such that*

$$x = \bigvee_{i\in I} x^i \ , \quad y = \bigwedge_{i\in I} y^i \ .$$

2.5. Lemma. $x \wedge y = a$ *and* $x \vee y = b$.

Proof. We apply the same steps as in proving the relations (4) and (5) in [6], Section 4 with the distinction that instead of infinite distributivity we apply 2.3 and the relation $L \in \mathcal{T}^1_\alpha$. □

The assertions of 4.3, 4.4 and 4.5 in [6] remain valid for our case (again, in the proof of 4.3 we have to use Lemma 2.3 above).

Now we can use the same argument as in Section 5 of [6] (instead of Lemma 4.2 of [5] we take Lemma 2.5 above). We apply the definitions of R and R' on L (cf. [5]) and we obtain

2.6. Lemma. *$L = (s^0)s^0_R \times s^0_{R'}$ and the relation*

$$s^0_R = \bigvee_{i \in I} L_i$$

is valid in the Boolean algebra $F(L, s^0)$.

By applying the well-known theorem of Smith and Tarski [12] (cf. also Sikorski [11], Chapter II, Theorem 20.1) we conclude from 2.6 that the following theorem holds.

2.7. Theorem. *Let α be an infinite cardinal and let $L \in \mathcal{T}^1_\alpha$. Then the Boolean algebra $F(L, s^0)$ is α-complete.*

3. On lattices belonging to $\mathcal{T}^2_\sigma$

If $\alpha = \aleph_0$, then instead of $\mathcal{T}^2_\alpha$ we write $\mathcal{T}^2_\sigma$.

Let L be a lattice belonging to $\mathcal{T}^2_\sigma$ and $s^0 \in L$. Suppose that for each $n \in N$ we have an internal direct product decomposition

$$L = (s^0)L_n \times L'_n \tag{1}$$

such that, whenever $n(1)$ and $n(2)$ are distinct positive integers, then

$$L_{n(1)} \cap L_{n(2)} = \{s^0\}. \tag{2}$$

We use analogous notation as in Section 2 with the distinction that we now have N instead of I.

In particular, the relation

$$s^0_R = \bigvee_{n \in N} L_n \tag{3}$$

is valid in the Boolean algebra $F(L, s^0)$; we have

$$L = (s^0)s^0_R \times (s^0_R)' \tag{4}$$

and, in view of the duality, (3) yields

$$(s^0_R)' = \bigcap_{n \in N} L'_n\ . \tag{5}$$

If a, b, x and y are as in 2.5, then we write

$$x = x(a, b), \quad y = y(a, b).$$

3.1. Lemma. *Let $x^0 \in L$. Then x^0_R is the set of all elements $z \in L$ such that there exist $u, v \in L$ with $x^0, z \in [u, v]$, $x(x^0, v) = v$ and $y(u, x^0) = u$.*

Proof. This is a consequence of the definition of R (cf. [5], Section 5). $\square$

3.2. Lemma. *Let m and n be distinct positive integers. Then $L_m \subsetneqq L'_n$.*

Proof. In view of (1) and according to 3.7 in [5] we have

$$L_m = (s^0)(L_m \cap L_n) \times (L_m \cap L'_n).$$

Thus according to (2),

$$L_m = (s^0)\{s^0\} \times (L_m \cap L'_n) = L_m \cap L'_n.$$

□

Since the element s^0 was arbitrarily chosen, we get

3.3. Corollary. *Let m, n be as in 3.2 and $x \in L$. Then*

$$x_{R_m} \subseteq x_{R'_n}.$$

3.4. Lemma. *Let $x^0 \in L$ and suppose that $(x^n)_{n\in N}$ is an indexed system of elements of L such that* (i) *this system is orthogonal over x^0, and* (ii) *$x^n \in x^0_{R_n}$ for each $n \in N$. Let $x = \bigvee_{n\in N} x^n$. Then for each $n \in N$, $xR'_n x^n$.*

Proof. Let $n \in N$. Since $L \in \mathcal{T}^2_\sigma$, there exists $t \in L$ with

$$t = \bigvee_{m\in N\setminus\{n\}} x^m .$$

According to 3.3, all elements x^m under consideration belong to $x^0_{R'_n}$. Thus t belongs to the set $x^0_{R'_n}$ as well. Clearly $x = x^n \vee t$. Then $xR'_n\ (x^n \vee x^0)$, whence $xR'_n x^n$. □

3.5. Lemma. *Let $(y^n)_{n\in N}$ be an indexed system of elements of L such that for each $n \in N$, $y^n \in L_n$. Then there exists $p \in s^0_R$ such that for each $n \in N$, $p(L_n) = y^n$.*

Proof. For each $n \in N$ we denote

$$y^n \vee s^0 = x^n, \quad y^n \wedge s^0 = z^n .$$

Then in view of (2), the indexed system $(x^n)_{n\in N}$ is orthogonal over s^0, and $(z^n)_{n\in N}$ is orthogonal under s^0. Hence there exist elements

$$x = \bigvee_{n\in N} x^n, \quad z = \bigwedge_{n\in N} z^n$$

in L. Thus xRs^0Rz, whence

$$[z, x] \subseteq s^0_R.$$

Also, $y^n \in s^0_R$ for each $n \in N$.

Let $n \in N$. There exists a uniquely determined element t^n in L such that

$$\{t^n\} = x^n_{R'_n} \cap z_{R_n} .$$

Then from the relation $z \leqq x^n$ we obtain that t^n belongs to the interval $[z, x^n]$ and hence $t^n \in s_R^0$. Put $p^n = t^n \wedge y^n$. We have $p^n \in [z, t^n]$, thus

$$p^n R_n z. \tag{6}$$

Therefore 3.2 yields that the indexed system $(p^n)_{n\in N}$ is orthogonal over z. Hence there exists

$$p = \bigvee_{n\in N} p^n$$

in L. Clearly $p \in [z, x] \subseteq s_R^0$. In view of 3.4, for each $n \in N$ we have

$$pR'_n p^n. \tag{7}$$

Since $x^n R'_n t^n$ we get

$$(x^n \wedge y^n) R'_n (t^n \wedge y^n),$$

thus $y^n R'_n p^n$. Hence in view of (7), $y^n R'_n p$. But $y^n \in L_n$ and hence $p(L_n) = y^n$. □

3.6. Lemma. *Let $x, y \in s_R^0$. Suppose that $x(L_n) = y(L_n)$ for each $n \in N$. Then $x = y$.*

Proof. Denote $a = x \wedge y$, $b = x \vee y$. Then $a(L_n) = b(L_n) = x(L_n)$ for each $n \in N$. It suffices to show that $a = b$.

Let $n \in N$. Put $a(L_n) = t$. Then $\{t\} = L_n \cap a_{R'_n}$. Hence $aR'_n t$ and similarly $bR'_n t$, which implies that $aR'_n b$.

We have $a, b \in s_R^0$. Then there exists an indexed system $(x^n)_{n\in N}$ which is orthogonal over a such that $aR_n x^n$ for each $n \in N$ and $\bigvee_{n\in N} x^n = b$ (cf. the definition of R in [6]).

From the relations $a \leqq x^n \leqq b$ and $aR'_n b$ we obtain $aR'_n x^n$, whence $a = x^n$ for each $n \in N$. Thus $b = a$. □

Consider the mapping

$$\varphi : s_R^0 \to \prod_{n\in N} L_n$$

defined by

$$\varphi(x) = (x(L_n))_{n\in N}$$

for each $x \in s_R^0$.

From the definition of φ we immediately obtain that φ is a homomorphism. In view of 3.5, φ is an epimorphism. According to 3.6, φ is a monomorphism. Hence φ is an isomorphism of s_R^0 onto $\prod_{n\in N} L_n$. All L_n are sublattices of s_R^0 containing the element s^0. If $x \in L_n$ for some $n \in N$, then $(\varphi(x))_n = x$ and $(\varphi(x))_m = s^0$ for $m \neq n$. Hence in view of (3) we have

3.7. Lemma. *Let* (1) *and* (3) *be valid. Then*

$$\bigvee_{n\in N} L_n = (s^0) \prod_{n\in N} L_n.$$

3.8. Theorem. *Let L_1 and L_2 be lattices belonging to $\mathcal{T}_\sigma^2$. Suppose that*

(i) *L_1 is isomorphic to some direct factor of L_2;*
(ii) *L_2 is isomorphic to some direct factor of L_1.*

Then L_1 is isomorphic to L_2.

Proof. It suffices to apply the same argument as in proving Theorem 2 of [6] (Section 6) with the distinction that instead of Lemma 6.3 from [6] we now use Lemma 3.7. □

Theorem 3.8 generalizes Theorem 2 of [6].

4. Examples

4.1. Let N be the set of all positive integers with the usual linear order and let A be a two-element lattice. Put $B = A \times N$, $L = B \cup \{\omega\}$, where $b < \omega$ for each $b \in B$. Then $L \in \mathcal{T}_\alpha^1 \cap \mathcal{T}_\alpha^2$ for each infinite cardinal α, but L fails to be infinitely distributive.

4.2. Let L be as in 4.1 and let L_1 be a sublattice of L such that $L_1 = L \setminus \{\omega\}$. Then $L_1 \in \mathcal{T}_\alpha^1 \cap \mathcal{T}_\alpha^2$, L_1 is infinitely distributive and fails to be σ-complete.

Now let us return to the conditions (b'_α), ($\mathrm{b}'_{\alpha d}$), (b''_α), ($\mathrm{b}''_{\alpha d}$), (c'_α) and ($\mathrm{c}'_{\alpha d}$). We denote the system of these condition by S. Let α be an arbitrary infinite cardinal.

It is obvious that (b'_α) $\Rightarrow$ (b''_α) and ($\mathrm{b}'_{\alpha d}$) $\Rightarrow$ ($\mathrm{b}''_{\alpha d}$).

4.3. Let F be the system of finite subsets of the set N; the system F is partially ordered by the set-theoretics inclusion. Then F satisfies all the conditions from S except (b'_α).

4.4. Let F be as in 4.3 and let F_1 be a lattice which is dual to F. Let α be an arbitrary infinite cardinal. Then F_1 satisfies all the conditions from S except the condition ($\mathrm{b}'_{\alpha d}$).

4.5. Let F be as in 4.3 and let N be the set of all positive integers with the natural linear order. The lattice dual to N will be denoted by N'. We may assume that $F \cap N' = \emptyset$. Put $L = F \cup N'$. The partial order in L is defined as follows: for each $x \in F$ and each $y \in N'$ we put $x < y$. If $x, y \in F$ or $x, y \in N'$, then the relation $x \geqq y$ has its original meaning (deduced from F or from N', respectively). The lattice L satisfies all conditions from S except (b'_α) and (b''_α).

4.6. Let L be as in 4.5 and let L_1 be a lattice dual to L. Then L_1 satisfies all conditions from S except ($\mathrm{b}'_{\alpha d}$) and ($\mathrm{b}''_{\alpha d}$).

4.7. Let L be as in 4.5. We denote by ω the greatest element of L. Further, let L_1 be the sublattice of L with the underlying set $F \cup \{\omega\}$. Then L_1 satisfies all the conditions of the system S except (c'_α).

4.8. Let L_1 be as in 4.7 and let L_2 be a lattice dual to L_1. Then L_2 satisfies all the conditions of S except ($\mathrm{c}'_{\alpha d}$).

References

.

[1] A. De Simone, D. Mundici, M. Navara, *A Cantor-Bernstein theorem for σ-complete MV-algebras*, Czechoslovak Math. J. (to appear).

[2] J. Jakubík, *Cantor-Bernstein theorem for lattice ordered groups*, Czechoslovak Math. J. **22** (1972), 159-175.

[3] J. Jakubík, *On complete lattice ordered groups with strong units*, Czechoslovak Math. J. **46** (1996), 221-230.

[4] J. Jakubík, *Convex isomorphisms of archimedean lattice ordered groups*, Mathware and Soft Computing **5** (1998), 49-56.

[5] J. Jakubík, *Cantor-Bernstein theorem for MV-algebras*, Czechoslovak Math. J. **49** (1999), 517–526.

[6] J. Jakubík, *Direct product decompositions of infinitely distributive lattices*, Math. Bohemica **125** (2000), 341-354.

[7] J. Jakubík, *On orthogonally σ-complete lattice ordered groups*, Czechoslovak Math. J. (to appear).

[8] J. Jakubík, *Convex mappings of archimedean MV-algebras*, Math. Slovaca, (submitted).

[9] J. Jakubík, *A theorem of Cantor-Bernstein type for orthogonally σ-complete pseudo MV-algebras*, (submitted).

[10] R. Sikorski, *A generalization of theorem of Banach and Cantor-Bernstein*, Coll. Math. **1** (1948), 140-144.

[11] R. Sikorski, *Boolean Algebras*, Second Edition, Springer Verlag, Berlin, 1964.

[12] E. C. Smith, jr., A. Tarski, *Higher degrees of distributivity and completeness in Boolean algebras*, Trans. Amer. Math. Soc. **84** (1957), 230-257.

[13] A. Tarski, *Cardinal Algebras*, New York, 1949.

Matematický ústav SAV, Grešákova 6, 040 01 Košice, Slovakia
E-mail address: `musavke@saske.sk`

This preprint series was founded in 1996. Its purpose is to present manuscripts of submitted or unpublished papers and reports of fellows of the Mathematical Institute of the Slovak Academy of Sciences in Bratislava and its Košice branch. The authors are fully responsible for the content of the preprints. Postal addresses:

Mathematical Institute
Slovak Academy of Sciences
Štefánikova 49
SK–814 73 Bratislava
SLOVAKIA

Tel./Fax (+421 7) 5249 7316

Mathematical Institute
Slovak Academy of Sciences
Grešákova 6
SK–040 01 Košice
SLOVAKIA

Tel./Fax (+421 95) 6228291

PostScript files of preprints are available at WWW address:
`http://www.mat.savba.sk/preprints`

Latest preprints of the series:

3/2002 *Ján Jakubík*: Generalized cardinal properties of lattices and lattice ordered groups

2/2002 *Anatolij Dvurečenskij*: Product effect algebras

1/2002 *Anatolij Dvurečenskij*: Perfect effect algebras are categorically equivalent with Abelian interpolation po-groups

15/2001 *Martin Kochol*: Equivalent versions of the 3-flow conjecture

14/2001 *Sylvia Pulmannová*: Compatibility and decompositions of effects

13/2001 *Anatolij Dvurečenskij, Thomas Vetterlein*: Archimedeaness and the MacNeille completion of pseudoeffect algebras and PO-groups

12/2001 *Anatolij Dvurečenskij, Thomas Vetterlein*: Infinitary lattice and Riesz properties for pseudoeffect algebras and PO-groups

11/2001 *Roman Frič*: Coproducts of D-posets and their application to probability

10/2001 *Roman Frič*: Remarks on sequential envelopes II

9/2001 *Martin Kochol*: Forbidden subgraphs for the smallest counterexample to the 5-flow conjecture

8/2001 *Martin Kochol*: Reduction of the 5-flow conjecture to cyclically 6-edge-connected snarks

7/2001 *Martin Kochol*: Decomposition formulas for the flow polynomial

6/2001 *Flaviano Battelli, Michal Fečkan*: Chaos arising near a topologically transversal homoclinic set

5/2001 *Antonio Di Nola, Anatolij Dvurečenskij*: MV-test spaces versus MV-algebras

4/2001 *Anatolij Dvurečenskij, Pekka Lahti, Kari Ylinen*: The uniqueness question in the multidimensional moment problem with applications to phase space observables

3/2001 *Anatolij Dvurečenskij, Pavel Pták*: Quantum logics with the Riesz interpolation property

2/2001 *Olga Nánásiová, Sylvia Pulmannová*: Abelian extensions of difference sets

1/2001 *Ján Jakubík*: Cantor-Bernstein theorem for lattices

23/2000 *Anatolij Dvurečenskij, Thomas Vetterlein*: Algebras in the positive cone of *po*-groups

22/2000 *Anatolij Dvurečenskij, Thomas Vetterlein*: Generalized pseudo-effect algebras

21/2000 *Martin Kochol*: Tension polynomials of graphs

20/2000 *Martin Kochol*: Sublinear defect principle in graph theory

(continued on inside back cover)

19/2000 *Martin Kochol*: Polynomials associated with nowhere-zero flows

18/2000 *Martin Kochol*: Superposition and constructions of graphs without nowhere-zero k-flows

17/2000 *Sylvia Pulmannová*: Divisible effect algebras and interval effect algebras

16/2000 *Gejza Jenča, Sylvia Pulmannová*: Quotients of partial abelian monoids and the Riesz decomposition property

15/2000 *Pekka Lahti, Sylvia Pulmannová*: Coexistence vs. functional coexistence of quantum observables

14/2000 *Beloslav Riečan*: Almost everywhere convergence in MV-algebras with product

13/2000 *Ján Jakubík*: Strong subdirect products of MV-algebras

12/2000 *Roman Frič*: Convergence and duality

11/2000 *Ján Jakubík*: Higher degrees of distributivity in MV-algebras

10/2000 *Beloslav Riečan*: On the probability on BL algebras

9/2000 *Beloslav Riečan*: On the Yager type entropy of dynamical systems

8/2000 *Anatolij Dvurečenskij*: States on pseudo MV-algebras

7/2000 *Anatolij Dvurečenskij, Thomas Vetterlein*: Pseudo-effect algebras. II. group representations

6/2000 *Anatolij Dvurečenskij, Thomas Vetterlein*: Pseudo-effect algebras. I. basic properties

5/2000 *Ján Jakubík*: On free MV-algebras

4/2000 *Anatolij Dvurečenskij*: Pseudo MV-algebras are intervals in l-groups

3/2000 *Anna Jenčová*: Geometry of quantum states: dual connections and divergence functions

2/2000 *Anatolij Dvurečenskij*: MV-observables and MV-algebras

1/2000 *Michal Fečkan*: Transversal bounded solutions for difference equations

21/1999 *Roman Frič*: Lukasiewicz tribes are absolutely sequentially closed bold algebras

20/1999 *Miroslav Repický*: Spaces not distinguishing convergences

19/1999 *Beloslav Riečan*: On the probability theory on MV algebras

18/1999 *Beloslav Riečan, Jozefína Petrovičová*: Kolmogorov-Sinai theorem for fuzzy T-dynamical systems

17/1999 *Beloslav Riečan*: On a Banach space valued observables

16/1999 *Michal Fečkan*: Criteria on the nonexistence of invariant submanifolds for dynamical systems

15/1999 *Anatolij Dvurečenskij*: On Loomis-Sikorski's theorem for MV-algebras and BCK-algebras

14/1999 *Otakar Grošek, Ladislav Satko, Karol Nemoga*: ZKDWCNDPDC...

13/1999 *Otakar Grošek, Karol Nemoga, Ladislav Satko*: A remark to Luby-Rackoff and Ueli M. Maurer pseudorandom generators

12/1999 *Beloslav Riečan*: On the sum of observables in product MV algebras

11/1999 *Peter Eliaš*: A hierarchy of thin sets related to the boundedness of trigonometric series

10/1999 *Miloslav Duchoň, Peter Maličký*: On Herglotz theorem in partially ordered vector spaces

9/1999 *Ján Jakubík*: Torsion classes of specker lattice ordered groups

8/1999 *Ján Jakubík*: Graph automorphisms and cells of lattices

7/1999 *Miloslav Duchoň, Peter Maličký*: Positive linear maps on trigonometric polynomials