Mathematical Institute
Slovak Academy of Sciences, Bratislava

Preprint 3/2009

Miloslav Duchoň
Peter Maličký

A Helly theorem for functions with values in metric spaces

December 28, 2009

PREPRINT SERIES

A HELLY THEOREM FOR FUNCTIONS WITH VALUES IN METRIC SPACES

Miloslav Duchoň - Peter Maličký

Abstract. We present a Helly type theorem for sequences of functions with values in metric spaces and apply it to representations of some mappings on the space of continuous functions. A generalization of the Riesz theorem is formulated and proved. More concretely, a representation of certain majored linear operators on the space of continuos functions into a complete metric space.

1. Introduction

It is well known that the following theorem is true, (Na, [6]), (BDS, [1]),(W, [7]).

Riesz representation theorem. *Every continuous linear functional L on the set of continuous functions f defined on $[0,1]$ has the form*

$$Lf = \int_0^1 f(s)dg(s) \tag{R}$$

with a function g of bounded variation on $[0,1]$.

This theorem has many extensions and generalizations with various proofs. One of the possible proofs is based on the Helly theorem [Na] and also on the moment problem theorem. It can be shown that the problem of determining the general continuous linear functionals on the set of continuous functions is equivalent to that of determining the set of all moment sequences. It is our purpose to extend this result for majored linear operators from continuous functions to Banach spaces.

Helly's theorem had been of some importance a long time above all in the probability theory in connection with a problem of moments of distributions. Recall that in this connection, real-valued nondecreasing functions f on the interval $[a,b]$ of the real line are considered and that the following facts are true.

(1) (*First Helly's theorem*) Given a uniformly bounded sequence (f_n) of real-valued nondecreasing functions, there exists a subsequence (f_{n_k}) of (f_n) converging to a real-valued nondecreasing function f on $[a,b]$.

1991 *Mathematics Subject Classification.* 28B99, 44A60.
Key words and phrases. metric space, vector space, vector function, bounded variation, majored operator.
Supported by grant agency Vega , 2/4137/07

(2) (*Second Helly's theorem*) Given a sequence (f_n) of real-valued nondecreasing functions on $[a, b]$, converging to a real-valued nondecreasing function f, then, for every continuous function g on $[a, b]$, we have

$$\lim_{n\to\infty} \int_a^b g(t)\, df_n(t) = \int_a^b g(t)\, df(t).$$

More generally there are true the following facts ([1],[4],[6],[[7]).

(1) (*First Helly's theorem*) Given the sequence (f_n) of complex-valued functions of uniformly bounded variation on $[a, b]$ such that for for some $x_0 \in [a, b]$ the sequence $(f_n(x_0))$ is bounded then there exists a subsequence (f_{n_k}) of (f_n) converging to a some function f of bounded variation on $[a, b]$.

(2) (*Second Helly's theorem*) Given a sequence (f_n) of functions of uniformly bounded variation on $[a, b]$, converging to a some function f of bounded variation, then, for every continuous function g on $[a, b]$, we have

$$\lim_{n\to\infty} \int_a^b g(t)\, df_n(t) = \int_a^b g(t)\, df(t)\,.$$

2. A Helly theorem in metric spaces

In this section we shall prove some facts concerning Helly's theorem in the context of metric spaces.

Definition. *Let D be a subset of the real line, (X, d) be metric space and $h : D \to X$ be a function. If the set of all sums $\sum_{i=1}^{n} d(h(t_{i-1}), h(t_i))$, where $(t_i)_{i=0}^{n}$ is an increasing sequence elements of D, is bounded then g is said to be a function of bounded variation on D. The corresponding least upper bound is a variation of function h on set D.*

Proposition. *Let D be a dense subset of the interval $[a, b]$, (X, d) be a complete metric space and $h : D \to X$ be a function of bounded variation on D.*

(i) *For any $t \in (a, b]$ (resp. $t \in [a, b)$) there exists limit $h(t-0)$ (resp. $h(t+0)$)*

(ii) *Function $f : (a, b] \to X$ (resp. $g : [a, b) \to X$) defined by $f(t) = h(t-0)$ (resp. $g(t) = h(t+0)$) are functions of bounded variation on $(a, b]$ (resp. $[a, b)$).*

(iii) *$f(t-0) = g(t-0) = h(t-0)$ for all $t \in (a, b]$ and $f(t+0) = g(t+0) = h(t+0)$ for all $t \in [a, b)$.*

(iv) *Function f (resp. g) is continuous at the point $t \in (a, b)$ if and only if $h(t-0) = h(t+0)$.*

(v) *For all $t \in (a, b)$ except possibly countable set $h(t-0) = h(t+0)$.*

Proof. Let $(t_i)_{i=0}^{\infty}$ be an increasing (resp. decreasing) sequence elements of D. Since (X, d) is a complete metric space, it is sufficient to show that $(h(t_i))_{i=0}^{\infty}$ is a Cauchy sequence. To see this, note that series $\sum_{i=1}^{\infty} d(h(t_{i-1}), h(t_i))$ is convergent and $d(h(t_n), h(t_m)) \le \sum_{i=n+1}^{m} d(h(t_{i-1}), h(t_i)) \le \sum_{i=n+1}^{\infty} d(h(t_{i-1}), h(t_i))$ for any $n < m$. It proves (i). Let $(t_i)_{i=0}^{n}$ be an increasing sequence elements of $(a, b]$. Take $\varepsilon > 0$

and a sequence $(s_i)_{i=0}^n$ of D such that $s_0 < t_0 < s_1 < t_1 < \ldots < t_{n-1} < s_n < t_n$ and $d(f(t_i), h(s_i)) < \frac{\varepsilon}{2n}$. Then $\sum_{i=1}^n d(f(t_{i-1}), f(t_i)) \leq \sum_{i=1}^n (d(f(t_{i-1}), h(s_{i-1})) + d(h(s_{i-1}), h(s_i)) + d(h(s_i), f(t_i))) \leq \sum_{i=1}^n d(h(s_{i-1}), h(s_i)) + \sum_{i=1}^n d(f(t_{i-1}), h(s_{i-1})) + \sum_{i=1}^n d(f(t_i), h(s_i)) < \varepsilon + \sum_{i=1}^n d(h(s_{i-1}), h(s_i))$. It means that f is of bounded variation. Bounded variation of g may be proved analogously. It proves (ii). Now we prove $f(t-0) = h(t-0)$ for $t \in (a, b]$. Let $(t_i)_{i=0}^\infty$ be an increasing sequence elements of $(a, b]$ with $\lim_{n\to\infty} t_n = t$. Take a sequence $(s_i)_{i=0}^\infty$ of D such that $s_0 < t_0 < s_1 < t_1 < \ldots < t_{n-1} < s_n < t_n < \ldots$ and $d(f(t_i), h(s_i)) < \frac{1}{n}$. Clearly $f(t-0) = \lim_{n\to\infty} f(t_n) = \lim_{n\to\infty} h(s_n) = h(t-0)$. Equalities $g(t-0) = h(t-0)$ for $t \in (a, b]$, $f(t+0) = h(t+0)$ and $g(t+0) = h(t+0)$ for $t \in [a, b)$ may be proved analogously. It proves (iii). Part (iv) follows from (iii). Let M be a variation of g on D and $n > 0$ be a fixed. Assume that there are m points $t_1, \ldots t_m$, where $m > nM$, such that inequality $d(f(t_i), h(t_i+0)) > \frac{1}{n}$ is satisfied for all $i = 1, \ldots, m$. There is $\varepsilon > 0$ such that $d(f(t_i), h(t_i+0)) > 2\varepsilon + \frac{1}{n}$ for all $i = 1, \ldots, m$. We may assume that $(t_i)_{i=1}^m$ is an increasing sequence. There are sequences $(s_i)_{i=1}^m$ and $(u_i)_{i=1}^m$ in D such that $s_1 < t_1 < u_1 < s_2 < t_2 < u_2 < \ldots, t_{m-1} < u_{m-1} < s_m < t_m < u_m$, $d(f(t_i), h(s_i)) < \varepsilon$ and $d(h(t_i+0), h(u_i)) < \varepsilon$ for all $i = 1, \ldots, m$. Now $\sum_{i=1}^m d(h(s_i), h(u_i)) > \sum_{i=1}^m (d(f(t_i), h(t_i+0)) - 2\varepsilon) >> \frac{m}{n} > M$ what is a contradiction. Therefore inequality $d(h(t-0), h(t+0)) > 0$ may be satisfied only for countably many $t \in (a, b)$.

Helly theorem. *Let (X, d) be a complete metric space and $(g_n)_{n\in N}$ a sequence of functions from $[a, b]$ into X such that a) the set $g_n(x)$ is relatively compact for any $x \in [a, b]$*

b) the functions $(g_n)_{n\in N}$ have uniformly bounded variations.

Then there exists a subsequence of the sequence $(g_n)_{n\in N}$ converging pointwise in X to a function $g : [a, b] \to X$ of bounded variation.

Proof. Let D be a countable set dense in $[a, b]$ and $b \in D$. By the diagonal procedure we obtain a subsequence $(h_n)_{n\in N}$ of $(g_n)_{n\in N}$ converging for every $t \in D$ to a function $h(t)$ which has bounded variation on D. Hence for every $t \in [a, b)$ there exists the limit $h(t+0)$. Put $g(t) = h(t+0)$, for $t \in [a, b)$ and $g(b) = h(b)$. Let $\delta > 0$ be given. We prove that

$$\limsup_{n\to\infty} d(h_n(t) - g(t)) \leq \delta$$

for every $t \in [a, b)$, except possible, finite number $m \leq 2M/\delta$, where M is the upper bound of variations of all g_n. So, assume that for some $m > 2M/\delta$ there are points $t_1 < t_2 < \ldots < t_m < b$ such that

$$\limsup_{n\to\infty} d(h_n(t_i) - g(t_i)) > \delta \text{ for all } i = 1, 2, \ldots, m\,.$$

There is a subsequence $(h_{n_k})_{k=1}^\infty$ such that

$$d(h_{n_k}(t_i) - g(t_i)) > \delta \text{ for all } i = 1, 2, \ldots, m \text{ and } k = 1, 2, \ldots$$

Since $g(t_i) = h(t_i + 0)$, there is a sequence $(u_i)_{i=1}^m$ in D such that

$$d(g(t_i), h(u_i)) < \frac{\delta}{4} \text{ for all } i = 1, 2, \ldots, m$$

and $t_1 < u_1 < t_2 < u_2 < \dots, t_{m-1} < u_{m-1} < t_m < u_m$. Let k be so large that

$$d(h_{n_k}(u_i), h(u_i)) \leq \delta/4, \text{ for all } i = 1, \dots, m\,.$$

Then

$$\sum_{i=1}^{m} d(h_{n_k}(t_i), h_{n_k}(u_i)) \geqq \sum_{i=1}^{m} (d(h_{n_k}(t_i), g(t_i)) - d(g(t_i), h(u_i)) - d(h(u_i), h_{n_k}(u_i)))$$

$$>> \sum_{i=1}^{m} (\delta - \delta/4 - \delta/4) = \frac{m\delta}{2} > M\,.$$

which is not possible. So,

$$\limsup_{n \to \infty} d(h_n(t) - g(t)) = 0$$

for every $t \in [a, b)$, except possible, countable set $A \subset [a, b)$. The last application of the diagonal procedure gives sequence convergent to $\widetilde{g}(t)$ on A. For some $t \in A$ it may be $\widetilde{g}(t) \neq g(t)$. Therefore g has to be redefined on A by $g(t) = \widetilde{g}(t)$.

Collorary of Helly theorem. *Let X be a complete linear metric space and $(g_n)_{n \in N}$ a sequence of functions from $[a, b]$ into X such that*

a) the set $\{g_n(x)\}$ is relatively compact for any $x \in [a, b]$

b) the functions $(g_n)_{n \in N}$ have uniformly bounded variations.

Then there exists a subsequence of the sequence $(g_n)_{n \in N}$ converging pointwise in X to a function $g : [a, b] \to X$ of bounded variation.

3. Integral representation theorem for majored linear operator

It is well known that every continuous linear form on the space of $C(I)$, $I = [0, 1]$, is presentable in the form

$$f \to \int_0^1 f(t)dq(t) \tag{3}$$

where q is a function of bounded variation.

From the preceding generalization of Helly theorem we obtain the result giving the representation of majored linear mapping ([4],Di). First we recall the definition of majored linear operator.

For each subset A of $[a, b]$, let $C([a, b], A)$ denote the space of continuous functions on $[a, b]$ vanishing outside A. Let X be a Banach space.If $F : C([a, b]) \to X$ is a linear mapping, define for each A,

$$|||F_A||| = \sup \sum_i ||F(f_i)||,$$

where the supremum is over all finite families $\{f_i\}$ in $C([a, b], A)$ with $\sum_i |f_i| \leq \chi_A(t)$ for all t in $[a, b]$. If $F : C([a, b]) \to X$ is a linear mapping, then (see [4], §19, p. 380, 383), the mapping F is called majored (also dominated, ([4]), if

$$|||F_A||| < \infty \tag{M}$$

for all A in $\mathcal{B}([a, b])$.

5. Riesz type theorem in some linear metric spaces

Let on the interval $[a,b]$ a function g of bounded variation with values in Banach X be given. This function may it possible to every continuous function $f(x)$ on $[a,b]$ to associate the element of X of the form

$$F(f)=\int_a^b f(x)dg(x). \tag{!}$$

The following properties are true. Let $f_1,f_2,f\in C[a,b]$. Then

$$F(f_1+f_2)=F(f_1)+F(f_2) \tag{a}$$

$$\|F(f)\|\le V_a^bM(f),\ M(f)=\max|f(x)|,\ V_a^b=Var_a^b(g) \tag{b}$$

Theorem (Riesz type). *Let on the set $C[a,b]$ the majored and compact linear mapping $F(f)$ with values in a Banach space X be given. Then there exists a function $g(x)$ of bounded variation with values X such that for every function $f(x)\in C[a,b]$ we have*

$$F(f)=\int_a^b f(x)dg(x) \tag{1}$$

Proof. It is enough to consider the case $a=1,b=1$, because the general case can be reduced to this case by means of a linear transformation of argument.

Put $\varphi_{n,k}=\binom{n}{k}x^k(1-x)^{n-k}$. It is easy to see that for every x it is true

$$\sum_{k=0}^n\varphi_{n,k}(x)=1.$$

Moreover, for $x\in[0,1]$ every member of this sum in nonnegative. Hence, if

$$a_k=\pm1,\ k=0,1,\dots,n,$$

then

$$|\sum_{k=0}^n a_k\varphi_{n,k}|\le 1. \tag{2}$$

Note that the considered linear operator $F(f)$ is defined for continuous on $[0,1]$ functions $f(x)$. According to definition of compact linear operator there exists a compact subset C of X such that

$$F(f)\in M(f)C,$$

or, equivalently mapping the unit sphere in $C[0,1]$ into C. From this and (2) we obtain

$$\sum_{k=0}^{n} a_k F(\varphi_{n,k}) \in C.$$

Further from the majority of operator F we obtain

$$\sum_{k=0}^{n} \|F(\varphi_{n,k})\| \leq \sup \sum_{i} \|F(f_i)\| \leq \||F_A\|| < \infty, \tag{3}$$

where the supremum is over all finite families $f_i \in C([a,b], A)$ with $\sum |f_i(t)| \leq \chi_A(t)$ for all $t \in [a,b]$, for all $A \in \mathcal{B}([a,b])$.

Let us define the step function $g_n(x)$ to put

$$g_n(0) = 0$$

$$g_n(x) = F(\varphi_{n,0}) \qquad (0 < x < \frac{1}{n})$$

$$g_n(x) = F(\varphi_{n,0}) + F(\varphi_{n,1}) \quad (\frac{1}{n} \leq x < \frac{2}{n})$$

$$\dots\dots\dots\dots\dots\dots\dots\dots\dots\dots$$

$$\dots\dots\dots\dots\dots\dots\dots\dots\dots\dots$$

$$g_n(x) = \sum_{k=0}^{n-1} F(\varphi_{n,k}) \quad (\frac{n-1}{n} \leq x < 1)$$

$$g_n(1) = \sum_{k=0}^{n} F(\varphi_{n,k})$$

By (3) the functions $g_n(x)$ and their total variations are bounded with the one number. Moreover, because by assumption $F(x)$ is compact operator on $C[0,1]$ into X , it takes unit sphere in $C[0,1]$ into a compact set C into X. Hence the set $\{g_n(x)\}, n = 1, 2, \dots$ is contained in C for all $x \in [0,1]$.

Hence on the base of Helly principle of choice from the sequence $\{g_n(x)\}$ it is possible to choose the subsequence $\{g_{n_i}(x)\}$ which converges in each point of $[0,1]$ to a function of the bounded variation.

If $f(x)$ is a continuous function on $[0,1]$, then on the base of (& 6, Th 3, [Na]) it can be shown that

$$\int_0^1 f(x) dg_n(x) = \sum_{k=0}^{n} f(\frac{k}{n}) F(\varphi_{n,k}),$$

from where

$$\int_0^1 f(x) dg_n(x) = F(B_n),$$

where

$$B_n(x) = \sum_{k=0}^{n} f(\frac{k}{n})\binom{n}{k} x^k(1-x)^{n-k},$$

is the Bernstein polynom for the function $f(x)$.

By theorem of S. N. Bernstein (&5, ch. IV, [Na])

$$M(B_n - f) \to 0, n \to \infty,$$

and by the definition of continuou linear operator we have

$$\|F(B_n) - F(f)\| = \|F(B_n - f)\| \leq KM(B_n - f).$$

This means that

$$F(B_n) \to F(f),$$

from where

$$\lim_{n\to\infty} \int_0^1 f(x)dg_n(x) = F(f).$$

But if $n \to \infty$, going through values $n_1, n_2, \dots$, then by Helly theorem (cf. also &7, [Na], and [2]), we obtain

$$\lim_{n\to\infty} \int_0^1 f(x)dg_n(x) = \int_0^1 f(x)dg(x).$$

Remark. We have derived here a representation theorem for majored operators using Helly theorem in linear metric spaces. To prove it by means of the Helly theorem from this paper we were able to do it for compact majored mappings. On the other hand, our approach is more "constructive" In ([4]) this is done for Banach spaces without such an assumption.

References

1. BARTLE, R. G., DUNFORD, N., SCHWARTZ, J., *Weak compactness and vector measures*, Canad. J. Math. **7** (1955), 289–305.
2. DEBIEVE, C., DUCHOŇ, M., DUHOUX, M., *A Helly theorem in the setting of Banach spaces*, Tatra Mount. Math. Publications **22** (2001), 105–114.
3. DEBIEVE, C., DUCHOŇ, M., DUHOUX, M., *A Helly's theorem in some Banach lattices*, Math. J. Toyama Univ. **23** (2000), 163–174.
4. DINCULEANU, N., *Vector Measures*, VEB, Berlin, 1966.
5. HAUSDORFF, F., *Momentprobleme für ein endliches Interval*, Math. Z. **16** (1923), 220–248.
6. NATANSON, I. P., *Theory of Functions of a Real Variable*, Frederick Ungar Publishing Co., New York, 1974.
7. WIDDER, D. V., *The Laplace Transform*, Princeton University Press, Princeton, 1946.

Mathematical Institute, Slovak Academy of Sciences, Štefánikova 49, SK-814 73 Bratislava, Slovakia

E-mail address: duchon@mat.savba.sk,

Faculty of natural sciences, University of Matej Bel, Podlavická, SK- Banská Bystrica, Slovakia

E-mail address: malicky@fpv.umb.sk

This preprint series was founded in 1996. Its purpose is to present manuscripts of submitted or unpublished papers and reports of fellows of the Mathematical Institute of the Slovak Academy of Sciences in Bratislava and its Košice branch. The authors are fully responsible for the content of the preprints. Postal addresses:

Mathematical Institute
Slovak Academy of Sciences
Štefánikova 49
SK–814 73 Bratislava
SLOVAKIA

Tel./Fax (+421 7) 397316

Mathematical Institute
Slovak Academy of Sciences
Grešákova 6
SK–040 01 Košice
SLOVAKIA

Tel./Fax (+421 95) 6228291

PostScript files of preprints are available at WWW address:
`http://www.savba.sk/sav/inst/math/intro.html`

Latest preprints of the series:

3/2009 *Miloslav Duchoň, Peter Maličký*: A Helly theorem for functions with values in metric spaces

2/2009 *Miloslav Duchoň*: Moments of vector measures and Pettis itegrable functions

1/2009 *Dilna, Natalyia, Fečkan, Michal*: Weakly nonlinear and symmetric periodic systems at resonance

4/2008 *Battelli, F., Fečkan, Michal*: On the chaotic behaviour of discontinuous systems

3/2008 *Dilna, Natalyia, Fečkan, Michal*: On the uniqueness and stability of symmetric and periodic solutions of weakly nonlinear ordinary differential equations

2/2008 *Fečkan, Michal, Gruendler, Joseph*: The existence of chaos in infinite dimensional non-resonant systems

1/2008 *Brandi, P., Ceppitelli, R., Holá, Ľ.*: Boundedly UC spaces and topologies on function spaces

10/2007 *Duchoň, Miloslav*: Moments of vector-valued functions and measures

9/2007 *Duchoň, Miloslav*: Moment problem for double fuzzy sequences

8/2007 *Fečkan, Michal*: Homoclinic bifurcations in discontinuous differential equations

7/2007 *Holá, Ľubica*: Spaces of densely continuous forms

6/2007 *Holá, Ľubica, McCoy, R. A.*: Relations approximated by continuous functions in the Vietoris topology

5/2007 *Holá, Ľubica, Holý, Dušan*: Minimal usco maps densely continuous forms and upper semicontinuous functions

4/2007 *Holá, Ľ., McCoy, R. A., Pelant, J.*: Approximations of relations by continuous functions

3/2007 *Duchoň, Miloslav*: Moment problem for double fuzzy sequences

2/2007 *Duchoň, Miloslav, Maličký, P.*: Fourier transform of a measure with values in a ordered vector space

1/2007 *Fečkan, Michal*: Note on arithmetic convolution equations

6/2006 *Papčo, Martin*: On effect algebras of fuzzy sets

5/2006 *Dvurečenskij, A.*: On n-perfect GMV-algebras

4/2006 *Di Nola, A., Dvurečenskij, A., Tsinakis, C.*: Perfect GMV-algebras

3/2006 *Fečkan, M., Rotho, V. M.*: Travelling waves in Hamiltonian system on 2D lattices with nearest neighbor interactions

(continued on inside back cover)

2/2006 *Pulmannová, S.*: Effect algebras with compressions

1/2006 *Fečkan, Michal*: Minimal periods of periodic solutions

19/2005 *Pulmannová, S., Vinceková, E.*: Remarks on the order for quantum observable

18/2005 *Duchoň, Miloslav, Maličký, P.*: A herglotz theorem in partially ordered vector spaces

17/2005 *Duchoň, M.*: On weakly compact operators

16/2005 *Wimmer, G., Witkovský, V.*: univariate linear calibration via replicated errors-in-variables model

15/2005 *Strauch, O., Blažeková, O.*: Distribution on the sequence $p_n/n \bmod 1$

14/2005 *Holá, Ľ., McCoy, R. A.*: relations approximated by continuous functions in the Vietoris topology

13/2005 *Fečkan, M., Ma, R., Thompson, B.*: Weakly coupled oscillators and topological degree

12/2005 *Dvurečenskij, A., Holland, W. Ch.*: Top varieties of generalized MV-algebras and unital lattice-ordered groups

11/2005 *Dvurečenskij, A., Rachůnek, J.*: On Riečan and Bosbach states for bounded non-commutative $R\ell$-monoids

10/2005 *Fečkan, M., Rothos, V.*: Kink-like periodic travelling waves for lattice equations with on-site and inter-site potentials

9/2005 *Buhagiar, D., Chetcuti, E., Dvurečenskij, A.*: Loomis-Sikorski representation of monotone σ-complete effect algebras

8/2005 *Battelli, F., Fečkan, M.*: Periodic solutions of symmetric elliptic singular systems: the higher codimension case

7/2005 *Armas, S., Mimura, M.*: A note on Serre's condition for orientability of fibre bundles

6/2005 *Dvurečenskij, A., Hyčko, M.*: Algebras on subintervals of BL-algebras, pseudo BL-algebras and bounded residuated ℓ-monoids

5/2005 *Dvurečenskij, A., Hyčko, M.*: On the existence of states for linear pseudo BL-algebras

4/2005 *Holá, Ľ., McCoy, R. A.*: Relations approximated by continuous functions

3/2005 *Holá, Ľ., Kupka, I.*: Strong convergence on compact sets and fixed points

2/2005 *Holá, Ľ., McCoy, R. A.*: cardinal invariants of the topology of uniform convergence on compact sets on the space of minimal usco maps

1/2005 *Aizicovici, S., Fečkan, M.*: Forced symmetric oscillations of evolution equations

4/2004 *Pulmannová, S.*: Spectral theorem for σ-complete lattice effect algebras

3/2004 *Pulmannová, S.*: A spectral theorem for $\sigma - MV$-algebras

2/2004 *Pulmannová, S.*: On fuzzy hidden variables

1/2004 *Battelli, Flaviano, Fečkan, Michal*: From transverse heteroclinic cycles to transverse homoclinic orbits

12/2003 *Chetcuti, E., Dvurečenskij, A.*: The state-space of the lattice of orthogonally closed subspaces

11/2003 *Frič, Roman*: Remarks on statistical maps and fuzzy (operational) random variables

10/2003 *Fečkan, Michal*: Periodically forced damped beam resting on nonlinear elastic bearings

9/2003 *Fečkan, Michal*: Forced oscillations of beams on elastic bearings

8/2003 *Chetcuti, Emmanuel, Dvurečenskij, Anatolij*: The existence of finitely additive states on orthogonaly closed subspaces of incomplete inner product spaces